



On the Solutions and Behavior of Rational Systems of Difference Equations

E. M. Elsayed^{1,2} and H. S. Gafel^{1,3}

ABSTRACT: In this paper, we obtain the expressions of the solutions of the following nonlinear systems of difference equations

$$x_{n+1} = \frac{x_{n-9}}{1 + x_{n-9}y_{n-4}}, \quad y_{n+1} = \frac{y_{n-9}}{\pm 1 \pm x_{n-4}y_{n-9}}, \quad n = 0, 1, \dots,$$

where the initial conditions $x_{-9}, x_{-8}, x_{-7}, x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0, y_{-9}, y_{-8}, y_{-7}, y_{-6}, y_{-5}, y_{-4}, y_{-3}, y_{-2}, y_{-1}, y_0$ are arbitrary non zero real numbers.

Key Words: Difference equations, recursive sequences, periodic solution, system of difference equations.

Contents

1	Introduction	1
2	On The System $x_{n+1} = \frac{x_{n-9}}{1+x_{n-9}y_{n-4}}, y_{n+1} = \frac{y_{n-9}}{1+x_{n-4}y_{n-9}}$	2
2.1	The Form of the Solutions of System (2)	3
2.2	Numerical Examples	6
3	On The System $x_{n+1} = \frac{x_{n-9}}{1+x_{n-9}y_{n-4}}, y_{n+1} = \frac{y_{n-9}}{1-x_{n-4}y_{n-9}}$	7
3.1	The Form of the Solutions of System (3)	8
3.2	Numerical Examples	9
4	On The Systems $x_{n+1} = \frac{x_{n-9}}{1+x_{n-9}y_{n-4}}, y_{n+1} = \frac{y_{n-9}}{-1 \pm x_{n-4}y_{n-9}}$	11
4.1	The Form of the Solutions of Systems (4.1) and (4.2)	11
4.2	Numerical Examples	13

1. Introduction

Many problems in Probability give rise to difference equations. Difference equations relate to differential equations as discrete mathematics relates to continuous mathematics. Anyone who has made a study of differential equations will know that even supposedly elementary examples can be hard to solve. By contrast, elementary difference equations are relatively easy to deal with. Aside from Probability, Computer Scientists take an interest in difference equations for a number of reasons. For example, difference equations frequently arise when determining the cost of an algorithm in big-O notation. Since difference equations are readily handled by program, a standard approach to solving a nasty differential equation is to convert it to an approximately equivalent difference equation see [1]- [30].

Many articles discuss difference equations systems, for example, Elsayed et al. [17] dealt with the solutions of the system of the difference equations

$$x_{n+1} = \frac{1}{x_{n-p}y_{n-p}}, \quad y_{n+1} = \frac{x_{n-p}y_{n-p}}{x_{n-q}y_{n-q}}.$$

Kurbanli et al. [22] discussed the behavior of positive solutions of the following system

$$x_{n+1} = \frac{x_{n-1}}{1 + x_{n-1}y_n}, \quad y_{n+1} = \frac{y_{n-1}}{1 + y_{n-1}x_n}.$$

El-Dessoky [7] considered the solutions of the following system

$$x_{n+1} = \frac{y_{n-1}y_{n-2}}{x_n(\pm 1 \pm y_{n-1}y_{n-2})}, \quad y_{n+1} = \frac{x_{n-1}x_{n-2}}{y_n(\pm 1 \pm x_{n-1}x_{n-2})}.$$

In [10] El-Dessoky and Elsayed investigated the form of the solution of the following systems of difference equations.

$$x_{n+1} = \frac{x_{n-2}}{\pm 1 \pm x_{n-2}z_{n-1}y_n}, \quad y_{n+1} = \frac{y_{n-2}}{\pm 1 \pm y_{n-2}x_{n-1}z_n}, \quad z_{n+1} = \frac{z_{n-2}}{\pm 1 \pm z_{n-2}y_{n-1}x_n}$$

Elabbasy et al. [6] devoted to investigate the local asymptotic stability, boundedness and periodic solutions of particular cases of the following general system of difference equations:

$$x_{n+1} = \frac{a_1y_{n-1} + a_2x_{n-3} + a_3}{a_4y_{n-1}x_{n-3} + a_5}, \quad y_{n+1} = \frac{b_1x_{n-1} + b_2y_{n-3} + b_3}{b_4x_{n-1}y_{n-3} + b_5}.$$

Elsayed et al. [15] obtained the expressions of the solutions of the following nonlinear systems of difference equations

$$x_{n+1} = \frac{y_n x_{n-2}}{y_n + y_{n-3}}, \quad y_{n+1} = \frac{x_n y_{n-2}}{\pm x_n \pm x_{n-3}}.$$

Khan et al. [20], studied the qualitative behavior of two systems of second-order rational difference equations.

Zhang et al. [30] discussed the boundedness, persistence, and global asymptotic stability of positive solution for a system of third-order rational difference equations

$$x_{n+1} = A + \frac{x_n}{y_{n-1}y_{n-2}}, \quad y_{n+1} = A + \frac{y_n}{x_{n-1}x_{n-2}}.$$

Elsayed and Ahmed [16] investigated the solutions and the periodicity of the following rational systems of difference equations of three-dimensional

$$x_{n+1} = \frac{y_n x_{n-2}}{x_{n-2} \pm z_{n-1}}, \quad y_{n+1} = \frac{z_n y_{n-2}}{y_{n-2} \pm x_{n-1}}, \quad z_{n+1} = \frac{x_n z_{n-2}}{z_{n-2} \pm y_{n-1}}.$$

Asiri et al. [1] studied the form of the solutions and the periodicity of the following third order systems of rational difference equations

$$x_{n+1} = \frac{y_{n-2}}{1 - y_{n-2}x_{n-1}y_n}, \quad y_{n+1} = \frac{x_{n-2}}{1 - x_{n-2}y_{n-1}x_n}.$$

The aim of this article is to obtain the expressions of the solutions of the following systems of difference equations

$$x_{n+1} = \frac{x_{n-9}}{1 + x_{n-9}y_{n-4}}, \quad y_{n+1} = \frac{y_{n-9}}{\pm 1 \pm x_{n-4}y_{n-9}}, \quad n = 0, 1, 2, \dots, \quad (1.1)$$

where the initial conditions $x_{-9}, x_{-8}, x_{-7}, x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0, y_{-9}, y_{-8}, y_{-7}, y_{-6}, y_{-5}, y_{-4}, y_{-3}, y_{-2}, y_{-1}, y_0$ are arbitrary non zero real numbers. Moreover, we obtain some numerical simulation to the equation are given to illustrate our results.

2. On The System $x_{n+1} = \frac{x_{n-9}}{1+x_{n-9}y_{n-4}}, y_{n+1} = \frac{y_{n-9}}{1+x_{n-4}y_{n-9}}$

In this section, we study the solution of the following system of difference equations

$$x_{n+1} = \frac{x_{n-9}}{1 + x_{n-9}y_{n-4}}, \quad y_{n+1} = \frac{y_{n-9}}{1 + x_{n-4}y_{n-9}}, \quad (2.1)$$

where the initial conditions $x_{-9}, x_{-8}, x_{-7}, x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0, y_{-9}, y_{-8}, y_{-7}, y_{-6}, y_{-5}, y_{-4}, y_{-3}, y_{-2}, y_{-1}, y_0$ are arbitrary non zero real numbers.

2.1. The Form of the Solutions of System (2)

The following theorem describes the form of the solutions of system (2).

Theorem 1. Suppose that $\{x_n, y_n\}$ are solutions of the system (2). Then for $n = 0, 1, 2, \dots$, we have the following formulas

$$\begin{aligned}
x_{10n-9} &= s \prod_{i=0}^{n-1} \frac{1+2ist}{1+(2i+1)st}, & x_{10n-8} &= k \prod_{i=0}^{n-1} \frac{1+2ikw}{1+(2i+1)kw}, \\
x_{10n-7} &= h \prod_{i=0}^{n-1} \frac{1+2ihv}{1+(2i+1)hv}, & x_{10n-6} &= g \prod_{i=0}^{n-1} \frac{1+2igu}{1+(2i+1)gu}, \\
x_{10n-5} &= f \prod_{i=0}^{n-1} \frac{1+2ifr}{1+(2i+1)fr}, & x_{10n-4} &= e \prod_{i=0}^{n-1} \frac{1+(2i+1)eq}{1+(2i+2)eq}, \\
x_{10n-3} &= d \prod_{i=0}^{n-1} \frac{1+(2i+1)dp}{1+(2i+2)dp}, & x_{10n-2} &= c \prod_{i=0}^{n-1} \frac{1+(2i+1)co}{1+(2i+2)co}, \\
x_{10n-1} &= b \prod_{i=0}^{n-1} \frac{1+(2i+1)Lb}{1+(2i+2)Lb}, & x_{10n} &= a \prod_{i=0}^{n-1} \frac{1+(2i+1)az}{1+(2i+2)az}, \\
y_{10n-9} &= q \prod_{i=0}^{n-1} \frac{1+2ieq}{1+(2i+1)eq}, & y_{10n-8} &= p \prod_{i=0}^{n-1} \frac{1+2idp}{1+(2i+1)dp}, \\
y_{10n-7} &= o \prod_{i=0}^{n-1} \frac{1+2ico}{1+(2i+1)co}, & y_{10n-6} &= L \prod_{i=0}^{n-1} \frac{1+2iLb}{1+(2i+1)Lb}, \\
y_{10n-5} &= z \prod_{i=0}^{n-1} \frac{1+2iaz}{1+(2i+1)az}, & y_{10n-4} &= t \prod_{i=0}^{n-1} \frac{1+(2i+1)st}{1+(2i+2)st}, \\
y_{10n-3} &= w \prod_{i=0}^{n-1} \frac{1+(2i+1)kw}{1+(2i+2)kw}, & y_{10n-2} &= v \prod_{i=0}^{n-1} \frac{1+(2i+1)hv}{1+(2i+2)hv}, \\
y_{10n-1} &= u \prod_{i=0}^{n-1} \frac{1+(2i+1)gu}{1+(2i+2)gu}, & y_{10n} &= r \prod_{i=0}^{n-1} \frac{1+(2i+1)fr}{1+(2i+2)fr},
\end{aligned}$$

where the initial conditions $x_{-9} = s$, $x_{-8} = k$, $x_{-7} = h$, $x_{-6} = g$, $x_{-5} = f$, $x_{-4} = e$, $x_{-3} = d$, $x_{-2} = c$, $x_{-1} = b$, $x_0 = a$, $y_{-9} = q$, $y_{-8} = p$, $y_{-7} = o$, $y_{-6} = L$, $y_{-5} = z$, $y_{-4} = t$, $y_{-3} = w$, $y_{-2} = v$, $y_{-1} = u$, $y_0 = r$.

Proof. For $n = 0$ the result holds. Suppose that the result holds for $n - 1$.

$$\begin{aligned}
x_{10n-19} &= s \prod_{i=0}^{n-2} \frac{1+2ist}{1+(2i+1)st}, & x_{10n-18} &= k \prod_{i=0}^{n-2} \frac{1+2ikw}{1+(2i+1)kw}, \\
x_{10n-17} &= h \prod_{i=0}^{n-2} \frac{1+2ihv}{1+(2i+1)hv}, & x_{10n-16} &= g \prod_{i=0}^{n-2} \frac{1+2igu}{1+(2i+1)gu}, \\
x_{10n-15} &= f \prod_{i=0}^{n-2} \frac{1+2ifr}{1+(2i+1)fr}, & x_{10n-14} &= e \prod_{i=0}^{n-2} \frac{1+(2i+1)eq}{1+(2i+2)eq}, \\
x_{10n-13} &= d \prod_{i=0}^{n-2} \frac{1+(2i+1)dp}{1+(2i+2)dp}, & x_{10n-12} &= c \prod_{i=0}^{n-2} \frac{1+(2i+1)co}{1+(2i+2)co},
\end{aligned}$$

$$x_{10n-11} = b \prod_{i=0}^{n-2} \frac{1 + (2i+1)Lb}{1 + (2i+2)Lb}, \quad x_{10n-10} = a \prod_{i=0}^{n-2} \frac{1 + (2i+1)az}{1 + (2i+2)az},$$

$$y_{10n-19} = q \prod_{i=0}^{n-2} \frac{1 + 2ieq}{1 + (2i+1)eq}, \quad y_{10n-18} = p \prod_{i=0}^{n-2} \frac{1 + 2idp}{1 + (2i+1)dp},$$

$$y_{10n-17} = o \prod_{i=0}^{n-2} \frac{1 + 2ico}{1 + (2i+1)co}, \quad y_{10n-16} = L \prod_{i=0}^{n-2} \frac{1 + 2iLb}{1 + (2i+1)Lb},$$

$$y_{10n-15} = z \prod_{i=0}^{n-2} \frac{1 + 2iaz}{1 + (2i+1)az}, \quad y_{10n-14} = t \prod_{i=0}^{n-2} \frac{1 + (2i+1)st}{1 + (2i+2)st},$$

$$y_{10n-13} = w \prod_{i=0}^{n-2} \frac{1 + (2i+1)kw}{1 + (2i+2)kw}, \quad y_{10n-12} = v \prod_{i=0}^{n-2} \frac{1 + (2i+1)hv}{1 + (2i+2)hv},$$

$$y_{10n-11} = u \prod_{i=0}^{n-2} \frac{1 + (2i+1)gu}{1 + (2i+2)gu}, \quad y_{10n-10} = r \prod_{i=0}^{n-2} \frac{1 + (2i+1)fr}{1 + (2i+2)fr},$$

from system (2) we can prove as follow

$$\begin{aligned} x_{10n-9} &= \frac{x_{10n-19}}{1 + x_{10n-19}y_{10n-14}} = \frac{s \prod_{i=0}^{n-2} \left(\frac{1+2ist}{1+(2i+1)st} \right)}{1 + st \prod_{i=0}^{n-2} \left(\frac{1+2ist}{1+(2i+1)st} \right) \left(\frac{1+(2i+1)st}{1+(2i+2)st} \right)} \\ &= \frac{s \prod_{i=0}^{n-2} \left(\frac{1+2ist}{1+(2i+1)st} \right)}{1 + st \prod_{i=0}^{n-2} \left(\frac{1+2ist}{1+(2i+2)st} \right)} = \frac{s \prod_{i=0}^{n-2} \left(\frac{1+2ist}{1+(2i+1)st} \right)}{1 + \left(\frac{st}{1+(2n-2)st} \right)} \left(\frac{1 + (2n-2)st}{1 + (2n-2)st} \right) \\ &= \frac{s(1 + (2n-2)st) \prod_{i=0}^{n-2} \left(\frac{1+2ist}{1+(2i+1)st} \right)}{1 + (2n-2)st + st} \\ &= \frac{s(1 + (2n-2)st)}{1 + (2n-1)st} \prod_{i=0}^{n-2} \left(\frac{1 + 2ist}{1 + (2i+1)st} \right) \\ &= s \prod_{i=0}^{n-1} \left(\frac{1 + 2ist}{1 + (2i+1)st} \right). \end{aligned}$$

Also, we get

$$\begin{aligned}
y_{10n-9} &= \frac{y_{10n-19}}{1 + y_{10n-19}x_{10n-14}} = \frac{q \prod_{i=0}^{n-2} \frac{1+2ieq}{1+(2i+1)eq}}{1 + q \prod_{i=0}^{n-2} \frac{1+2ieq}{1+(2i+1)eq} e \prod_{i=0}^{n-2} \frac{1+(2i+1)eq}{1+(2i+2)eq}} \\
&= \frac{q \prod_{i=0}^{n-2} \frac{1+2ieq}{1+(2i+1)eq}}{1 + eq \prod_{i=0}^{n-2} \frac{1+2ieq}{1+(2i+2)eq}} = \frac{q \prod_{i=0}^{n-2} \frac{1+2ieq}{1+(2i+1)eq}}{1 + \frac{eq}{1+(2n-2)eq}} = \frac{q \prod_{i=0}^{n-2} \frac{1+2ieq}{1+(2i+1)eq}}{\frac{1+(2n-1)eq}{1+(2n-2)eq}} \\
&= q \prod_{i=0}^{n-2} \frac{1+2ieq}{1+(2i+1)eq} \left(\frac{1+(2n-2)eq}{1+(2n-1)eq} \right) = q \prod_{i=0}^{n-1} \frac{1+2ieq}{1+(2i+1)eq}.
\end{aligned}$$

The other relations can be proved by similar way. This completes the proof.

Lemma 1. The equilibrium points of system (2) are $(0, \alpha)$ and $(\gamma, 0)$ where $\alpha, \gamma \in [0, \infty)$.

Proof. For the equilibrium points of system (2), we can write

$$\bar{x} = \frac{\bar{x}}{1 + \bar{x}\bar{y}}, \quad \bar{y} = \frac{\bar{y}}{1 + \bar{x}\bar{y}}.$$

Then

$$\bar{x}(1 + \bar{x}\bar{y}) = \bar{x}, \quad \bar{y}(1 + \bar{x}\bar{y}) = \bar{y},$$

we have

$$\bar{x}(1 + \bar{x}\bar{y} - 1) = 0, \quad \bar{y}(1 + \bar{x}\bar{y} - 1) = 0.$$

Therefore every $(0, \alpha)$ and $(\gamma, 0)$ are solutions. Thus the equilibrium points of system (2) are $(0, \alpha)$ and $(\gamma, 0)$.

Lemma 2. Every positive solution of the system (2) is bounded and convergent.

Proof. Let $\{x_n, y_n\}$ be a positive solution of system (2). It follows from system (2) that

$$x_{n+1} = \frac{x_{n-9}}{1 + x_{n-9}y_{n-4}} < x_{n-9},$$

and

$$y_{n+1} = \frac{y_{n-9}}{1 + y_{n-9}x_{n-4}} < y_{n-9}.$$

Then the subsequences $\{x_{10n-9}\}_{n=0}^{\infty}$, $\{x_{10n-8}\}_{n=0}^{\infty}$, $\{x_{10n-7}\}_{n=0}^{\infty}$, $\{x_{10n-6}\}_{n=0}^{\infty}$, $\{x_{10n-5}\}_{n=0}^{\infty}$, $\{x_{10n-4}\}_{n=0}^{\infty}$, $\{x_{10n-3}\}_{n=0}^{\infty}$, $\{x_{10n-2}\}_{n=0}^{\infty}$, $\{x_{10n-1}\}_{n=0}^{\infty}$ and $\{x_{10n}\}_{n=0}^{\infty}$ are decreasing and so are bounded from above by

$$M = \max \{x_{-9}, x_{-8}, x_{-7}, x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0\}.$$

Similarly the subsequences $\{y_{10n-9}\}_{n=0}^{\infty}$, $\{y_{10n-8}\}_{n=0}^{\infty}$, $\{y_{10n-7}\}_{n=0}^{\infty}$, $\{y_{10n-6}\}_{n=0}^{\infty}$, $\{y_{10n-5}\}_{n=0}^{\infty}$, $\{y_{10n-4}\}_{n=0}^{\infty}$, $\{y_{10n-3}\}_{n=0}^{\infty}$, $\{y_{10n-2}\}_{n=0}^{\infty}$, $\{y_{10n-1}\}_{n=0}^{\infty}$ and $\{y_{10n}\}_{n=0}^{\infty}$ are decreasing and so are bounded from above by

$$N = \max \{y_{-9}, y_{-8}, y_{-7}, y_{-6}, y_{-5}, y_{-4}, y_{-3}, y_{-2}, y_{-1}, y_0\}.$$

Lemma 3. If $a, b, c, d, e, f, g, h, k, s, o, p, q, r, s, L, t, u, v, w$ and z arbitrary real numbers and let $\{x_n, y_n\}$ be a solution of system (2) then the following statements are true:

(i) If $a = 0$, $z \neq 0$, (or $z = 0$, $a \neq 0$), then $x_{10n} = 0$ and $y_{10n-5} = z$ (or $x_{10n} = a$ and $y_{10n-5} = 0$).

(ii) If $b = 0$, $L \neq 0$, (or $L = 0$, $b \neq 0$), then $x_{10n-1} = 0$ and $y_{10n-6} = L$ (or $x_{10n-1} = b$ and $y_{10n-6} = 0$).

(iii) If $c = 0$, $o \neq 0$, (or $o = 0$, $c \neq 0$), then $x_{10n-2} = 0$ and $y_{10n-7} = o$ (or $x_{10n-2} = c$ and $y_{10n-7} = 0$).

(iv) If $d = 0$, $p \neq 0$, (or $p = 0$, $d \neq 0$), then $x_{10n-3} = 0$ and $y_{10n-8} = p$ (or $x_{10n-3} = d$ and $y_{10n-8} = 0$).

(v) If $e = 0$, $q \neq 0$, (or $q = 0$, $e \neq 0$), then $x_{10n-4} = 0$ and $y_{10n-9} = q$ (or $x_{10n-4} = e$ and $y_{10n-9} = 0$).

(vi) If $f = 0$, $r \neq 0$, (or $r = 0$, $f \neq 0$), then $x_{10n-5} = 0$ and $y_{10n} = r$ (or $x_{10n-5} = f$ and $y_{10n} = 0$).

(vii) If $g = 0$, $u \neq 0$, (or $u = 0$, $g \neq 0$), then $x_{10n-6} = 0$ and $y_{10n-1} = u$ (or $x_{10n-6} = g$ and $y_{10n-1} = 0$).

(viii) If $h = 0$, $v \neq 0$, (or $v = 0$, $h \neq 0$), then $x_{10n-7} = 0$ and $y_{10n-2} = v$ (or $x_{10n-7} = h$ and $y_{10n-2} = 0$).

(ix) If $k = 0$, $w \neq 0$, (or $w = 0$, $k \neq 0$), then $x_{8n-1} = 0$ and $y_{10n-3} = w$ (or $x_{8n-1} = k$ and $y_{10n-3} = 0$).


(x) If $s = 0$, $t \neq 0$, (or $t = 0$, $s \neq 0$), then $x_{10n-9} = 0$ and $y_{10n-4} = t$ (or $x_{10n-9} = s$ and $y_{10n-4} = 0$).

Proof. The proof follows from the form of the solutions of system (1).


2.2. Numerical Examples

In order to illustrate the results of the previous sections and to support our theoretical discussions, we consider several interesting numerical examples in this section.


Example 1. For the initial conditions $x_{-9} = -1.5$, $x_{-8} = -6.2$, $x_{-7} = 0.6$, $x_{-6} = 1.3$, $x_{-5} = 12$, $x_{-4} = -0.6$, $x_{-3} = 0.9$, $x_{-2} = 0.2$, $x_{-1} = 0.6$, $x_0 = 8.3$, $y_{-9} = 0.7$, $y_{-8} = 1.8$, $y_{-7} = 0.3$, $y_{-6} = 1.1$, $y_{-5} = -1.4$, $y_{-4} = -2.5$, $y_{-3} = 1.6$, $y_{-2} = 9$, $y_{-1} = 0.3$, $y_0 = 0.9$ when we take the system (2). (See Fig. 1).



Example 2. We consider interesting numerical example for the difference equations system (2) with the initial conditions $x_{-9} = 1.5$, $x_{-8} = 6.2$, $x_{-7} = 6$, $x_{-6} = 13$, $x_{-5} = 12$, $x_{-4} = 0.6$, $x_{-3} = 9$, $x_{-2} = 2$, $x_{-1} = 0.6$, $x_0 = 8.3$, $y_{-9} = 0.7$, $y_{-8} = 1.8$, $y_{-7} = 0.3$, $y_{-6} = 11$, $y_{-5} = 14$, $y_{-4} = 25$, $y_{-3} = 16$, $y_{-2} = 9$, $y_{-1} = 0.3$, $y_0 = 2.9$. (See Fig. 2).



Example 3. Consider the difference system (2) with the initial conditions $x_{-9} = 1.5, x_{-8} = 0, x_{-7} = 6, x_{-6} = 13, x_{-5} = 0, x_{-4} = 0, x_{-3} = 9, x_{-2} = 0, x_{-1} = 0.6, x_0 = 0, y_{-9} = 0, y_{-8} = 1.8, y_{-7} = 0.3, y_{-6} = 0, y_{-5} = 14, y_{-4} = 0, y_{-3} = 16, y_{-2} = 0, y_{-1} = 0.3, y_0 = 2.9$. (See Fig. 3).



3. On The System $x_{n+1} = \frac{x_{n-9}}{1+x_{n-9}y_{n-4}}, y_{n+1} = \frac{y_{n-9}}{1-x_{n-4}y_{n-9}}$

In this section, we obtain the form of the solution of the following system of difference equations

$$x_{n+1} = \frac{x_{n-9}}{1+x_{n-9}y_{n-4}}, \quad y_{n+1} = \frac{y_{n-9}}{1-x_{n-4}y_{n-9}}, \tag{3.1}$$

where the initial conditions $x_{-9}, x_{-8}, x_{-7}, x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0, y_{-9}, y_{-8}, y_{-7},$

$y_{-6}, y_{-5}, y_{-4}, y_{-4}, y_{-3}, y_{-2}, y_{-1}, y_0$ are arbitrary non zero real numbers with $x_{-9}y_{-4}, x_{-8}y_{-3}, x_{-7}y_{-2}, x_{-6}y_{-1}, x_{-5}y_0 \neq -1$ and $x_{-4}y_{-9}, x_{-3}y_{-8}, x_{-2}y_{-7}, x_{-1}y_{-6}, x_0y_{-5} \neq 1$.

3.1. The Form of the Solutions of System (3)

Theorem 2. Let $\{x_n, y_n\}$ are solutions of system (3). Then for $n = 0, 1, 2, \dots$, we have

$$\begin{aligned} x_{10n-9} &= \frac{s}{(1+st)^n}, \quad x_{10n-8} = \frac{k}{(1+kw)^n}, \quad x_{10n-7} = \frac{h}{(1+hv)^n}, \quad x_{10n-6} = \frac{g}{(1+gu)^n}, \\ x_{10n-5} &= \frac{f}{(1+fr)^n}, \quad x_{10n-4} = (-1)^n e(eq-1)^n, \quad x_{10n-3} = (-1)^n d(dp-1)^n, \\ x_{10n-2} &= (-1)^n c(co-1)^n, \quad x_{10n-1} = (-1)^n b(Lb-1)^n, \quad x_{10n} = (-1)^n a(az-1)^n, \\ \\ y_{10n-9} &= \frac{(-1)^n q}{(eq-1)^n}, \quad y_{10n-8} = \frac{(-1)^n p}{(dp-1)^n}, \quad y_{10n-7} = \frac{(-1)^n o}{(co-1)^n}, \quad y_{10n-6} = \frac{(-1)^n L}{(Lb-1)^n}, \\ y_{10n-5} &= \frac{(-1)^n z}{(az-1)^n}, \quad y_{10n-4} = t(st+1)^n, \quad y_{10n-3} = w(kw+1)^n, \\ y_{10n-2} &= v(hv+1)^n, \quad y_{10n-1} = u(gu+1)^n, \quad y_{10n} = r(fr+1)^n. \end{aligned}$$

Proof. For $n = 0$ the result holds. Suppose that the result holds for $n - 1$

$$\begin{aligned} x_{10n-19} &= \frac{s}{(1+st)^{n-1}}, \quad x_{10n-18} = \frac{k}{(1+kw)^{n-1}}, \quad x_{10n-17} = \frac{h}{(1+hv)^{n-1}}, \\ x_{10n-16} &= \frac{g}{(1+gu)^{n-1}}, \quad x_{10n-15} = \frac{f}{(1+fr)^{n-1}}, \quad x_{10n-14} = (-1)^{n-1} e(eq-1)^{n-1}, \\ x_{10n-13} &= (-1)^{n-1} d(dp-1)^{n-1}, \quad x_{10n-12} = (-1)^{n-1} c(co-1)^{n-1}, \\ x_{10n-11} &= (-1)^{n-1} b(Lb-1)^{n-1}, \quad x_{10n-10} = (-1)^{n-1} a(az-1)^{n-1}, \\ \\ y_{10n-19} &= \frac{(-1)^{n-1} q}{(eq-1)^{n-1}}, \quad y_{10n-18} = \frac{(-1)^{n-1} p}{(dp-1)^{n-1}}, \quad y_{10n-17} = \frac{(-1)^{n-1} o}{(co-1)^{n-1}}, \\ y_{10n-16} &= \frac{(-1)^{n-1} L}{(Lb-1)^{n-1}}, \quad y_{10n-15} = \frac{(-1)^{n-1} z}{(az-1)^{n-1}}, \quad y_{10n-14} = t(st+1)^{n-1}, \\ y_{10n-13} &= w(kw+1)^{n-1}, \quad y_{10n-12} = v(hv+1)^{n-1}, \\ y_{10n-11} &= u(gu+1)^{n-1}, \quad y_{10n-10} = r(fr+1)^{n-1}, \end{aligned}$$

from system (3) we can see that

$$\begin{aligned} x_{10n-9} &= \frac{x_{10n-19}}{1 + x_{10n-19}y_{10n-14}} = \frac{\frac{s}{(1+st)^{n-1}}}{1 + \left(\frac{s}{(1+st)^{n-1}}\right)t(st+1)^{n-1}} \\ &= \frac{\frac{s}{(1+st)^{n-1}}}{1 + st} = \frac{s}{(1+st)^n}. \end{aligned}$$

Also, we get

$$\begin{aligned}
x_{10n-8} &= \frac{x_{10n-18}}{1 + x_{10n-18}y_{10n-13}} = \frac{\frac{k}{(1+kw)^{n-1}}}{1 + \left(\frac{k}{(1+kw)^{n-1}}w(kw+1)^{n-1}\right)} = \frac{k}{(1+kw)^n}, \\
x_{10n-3} &= \frac{x_{10n-13}}{1 + x_{10n-13}y_{10n-8}} = \frac{(-1)^{n-1}d(dp-1)^{n-1}}{1 + \frac{(-1)^n p}{(dp-1)^n}(-1)^{n-1}d(dp-1)^{n-1}} \\
&= \frac{(-1)^n d(dp-1)^n}{-dp+1+dp} = (-1)^n d(dp-1)^n, \\
y_{10n-6} &= \frac{y_{10n-16}}{1 - y_{10n-16}x_{10n-11}} = \frac{\frac{(-1)^{n-1}L}{(Lb-1)^{n-1}}}{1 - \frac{(-1)^{n-1}L}{(Lb-1)^{n-1}}(-1)^{n-1}b(Lb-1)^{n-1}} \\
&= \frac{\frac{(-1)^{n-1}L}{(Lb-1)^{n-1}}}{1 - (-1)^{2n-1}Lb} = \frac{(-1)^n L}{(Lb-1)^n}.
\end{aligned}$$

By similar way we can prove the other relations. This completes the proof.

Lemma 4. Assume that $\{x_n, y_n\}$ be a solution of system (3) with $x_{-9}y_{-4}, x_{-8}y_{-3}, x_{-7}y_{-2}, x_{-6}y_{-1}, x_{-5}y_0 = -2$ and $x_{-4}y_{-9}, x_{-3}y_{-8}, x_{-2}y_{-7}, x_{-1}y_{-6}, x_0y_{-5} = 2$, then $\{x_n\}$ and $\{y_n\}$ is periodic with period twenty.

Proof. From the form of the solutions of system (3) we see when $x_{-9}y_{-4}, x_{-8}y_{-3}, x_{-7}y_{-2}, x_{-6}y_{-1}, x_{-5}y_0 = -2$ and $x_{-4}y_{-9}, x_{-3}y_{-8}, x_{-2}y_{-7}, x_{-1}y_{-6}, x_0y_{-5} = 2$ that

$$\begin{aligned}
x_{10n-9} &= \frac{s}{(-1)^n}, x_{10n-8} = \frac{k}{(-1)^n}, x_{10n-7} = \frac{h}{(-1)^n}, x_{10n-6} = \frac{g}{(-1)^n}, x_{10n-5} = \frac{f}{(-1)^n}, \\
x_{10n-4} &= (-1)^n e, x_{10n-3} = (-1)^n d, x_{10n-2} = (-1)^n c, x_{10n-1} = (-1)^n b, x_{10n} = (-1)^n a,
\end{aligned}$$

$$\begin{aligned}
y_{10n-9} &= (-1)^n q, y_{10n-8} = (-1)^n p, y_{10n-7} = (-1)^n o, y_{10n-6} = (-1)^n L, y_{10n-5} = (-1)^n z, \\
y_{10n-4} &= t(-1)^n, y_{10n-3} = w(-1)^n, y_{10n-2} = v(-1)^n, y_{10n-1} = u(-1)^n, y_{10n} = r(-1)^n.
\end{aligned}$$

and so it is periodic with period twenty. This completes the proof.

Lemma 5. Let $\{x_n, y_n\}$ be a positive solution of system (3), then $\{x_n\}$ is bounded and converges to zero.

Proof. It follows from system (3) that

$$x_{n+1} = \frac{x_{n-9}}{1 + x_{n-9}y_{n-4}} < x_{n-9}.$$


Then the subsequences $\{x_{10n-9}\}_{n=0}^{\infty}, \{x_{10n-8}\}_{n=0}^{\infty}, \{x_{10n-7}\}_{n=0}^{\infty}, \{x_{10n-6}\}_{n=0}^{\infty}, \{x_{10n-5}\}_{n=0}^{\infty}, \{x_{10n-4}\}_{n=0}^{\infty}, \{x_{10n-3}\}_{n=0}^{\infty}, \{x_{10n-2}\}_{n=0}^{\infty}, \{x_{10n-1}\}_{n=0}^{\infty}$ and $\{x_{10n}\}_{n=0}^{\infty}$ are decreasing and so are bounded from above by

$$M = \max \{x_{-9}, x_{-8}, x_{-7}, x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0\}.$$


3.2. Numerical Examples

Here we plot some examples to illustrate the results of the previous sections.


Example 4. See Figure 4 when we suppose the initial conditions for system (3) as follows: $x_{-9} = 0.25, x_{-8} = 0.8, x_{-7} = 0.3, x_{-6} = 0.13, x_{-5} = 0.9, x_{-4} = 0.11, x_{-3} = 0.58, x_{-2} = 0.7, x_{-1} = 0.6, x_0 = 0.4, y_{-9} = 1.2, y_{-8} = .08, y_{-7} = 0.3, y_{-6} = 0.5, y_{-5} = 0.34, y_{-4} = 0.11, y_{-3} = 0.16, y_{-2} = 0.565, y_{-1} = 0.311, y_0 = 0.9.$



Example 5. We take a numerical example for the system (3) with the initial conditions $x_{-9} = 0.15$, $x_{-8} = -0.8$, $x_{-7} = 0.3$, $x_{-6} = -0.13$, $x_{-5} = 0.9$, $x_{-4} = -0.11$, $x_{-3} = -0.8$, $x_{-2} = 0.1$, $x_{-1} = 0.2$, $x_0 = -0.4$, $y_{-9} = 0.2$, $y_{-8} = -0.08$, $y_{-7} = 0.78$, $y_{-6} = 0.085$, $y_{-5} = 0.34$, $y_{-4} = 0.11$, $y_{-3} = 0.16$, $y_{-2} = -0.65$, $y_{-1} = 0.311$, $y_0 = 0.9$. (See Fig. 5).



Example 6. Figure 6 shows the periodic solutions with period twenty for system (3) when $x_{-9} = 2/11$, $x_{-8} = -1/3$, $x_{-7} = -4$, $x_{-6} = -20$, $x_{-5} = 0.2$, $x_{-4} = 0.4$, $x_{-3} = -0.25$, $x_{-2} = 2/9$, $x_{-1} = 20$, $x_0 = 1/7$, $y_{-9} = 5$, $y_{-8} = -8$, $y_{-7} = 9$, $y_{-6} = 0.1$, $y_{-5} = 14$, $y_{-4} = -11$, $y_{-3} = 6$, $y_{-2} = 0.5$, $y_{-1} = 0.1$, $y_0 = -10$.



4. On The Systems $x_{n+1} = \frac{x_{n-9}}{1+x_{n-9}y_{n-4}}$, $y_{n+1} = \frac{y_{n-9}}{-1\pm x_{n-4}y_{n-9}}$

Here, we investigate the expressions of the solutions of the following systems of difference equations

$$x_{n+1} = \frac{x_{n-9}}{1+x_{n-9}y_{n-4}}, \quad y_{n+1} = \frac{y_{n-9}}{-1+x_{n-4}y_{n-9}}, \quad (4.1)$$

$$x_{n+1} = \frac{x_{n-9}}{1+x_{n-9}y_{n-4}}, \quad y_{n+1} = \frac{y_{n-9}}{-1-x_{n-4}y_{n-9}}, \quad (4.2)$$

where the initial conditions $x_{-9}, x_{-8}, x_{-7}, x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0, y_{-9}, y_{-8}, y_{-7}, y_{-6}, y_{-5}, y_{-4}, y_{-3}, y_{-2}, y_{-1}, y_0$ are arbitrary non zero real numbers.

4.1. The Form of the Solutions of Systems (4.1) and (4.2)

Theorem 3. Assume that $\{x_n, y_n\}$ are solutions of the system (4.1) where the initial conditions satisfy that $x_{-9}y_{-4}, x_{-8}y_{-3}, x_{-7}y_{-2}, x_{-6}y_{-1}, x_{-5}y_0 \neq \pm 1$ and $x_{-4}y_{-9}, x_{-3}y_{-8}, x_{-2}y_{-7}, x_{-1}y_{-6}, x_0y_{-5} \neq 1, \neq \frac{1}{2}$. Then we have the following expressions for $n = 0, 1, 2, \dots$,

$$\begin{aligned} x_{20n-9} &= \frac{(-1)^n s}{(s^2 t^2 - 1)^n}, & x_{20n-8} &= \frac{(-1)^n k}{(k^2 w^2 - 1)^n}, & x_{20n-7} &= \frac{(-1)^n h}{(h^2 v^2 - 1)^n}, \\ x_{20n-6} &= \frac{(-1)^n g}{(g^2 u^2 - 1)^n}, & x_{20n-5} &= \frac{(-1)^n f}{(f^2 r^2 - 1)^n}, & x_{20n-4} &= \frac{(-1)^n e(eq - 1)^{2n}}{(2eq - 1)^n}, \\ x_{20n-3} &= \frac{(-1)^n d(dp - 1)^{2n}}{(2dp - 1)^n}, & x_{20n-2} &= \frac{(-1)^n c(co - 1)^{2n}}{(2co - 1)^n}, & x_{20n-1} &= \frac{(-1)^n b(Lb - 1)^{2n}}{(2Lb - 1)^n}, \\ x_{20n} &= \frac{(-1)^n a(az - 1)^{2n}}{(2az - 1)^n}, & x_{20n+1} &= \frac{(-1)^n s}{(st + 1)(s^2 t^2 - 1)^n}, & x_{20n+2} &= \frac{(-1)^n k}{(1 + kw)(k^2 w^2 - 1)^n}, \\ x_{20n+3} &= \frac{(-1)^n h}{(1 + hv)(h^2 v^2 - 1)^n}, & x_{20n+4} &= \frac{(-1)^n g}{(1 + gu)(g^2 u^2 - 1)^n}, & x_{20n+5} &= \frac{(-1)^n f}{(1 + fr)(f^2 r^2 - 1)^n}, \\ x_{20n+6} &= \frac{(-1)^n e(eq - 1)^{2n+1}}{(2eq - 1)^{n+1}}, & x_{20n+7} &= \frac{(-1)^n d(dp - 1)^{2n+1}}{(2dp - 1)^{n+1}}, & x_{20n+8} &= \frac{(-1)^n c(co - 1)^{2n+1}}{(2co - 1)^{n+1}}, \\ x_{20n+9} &= \frac{(-1)^n b(Lb - 1)^{2n+1}}{(2Lb - 1)^{n+1}}, & x_{20n+10} &= \frac{(-1)^n a(az - 1)^{2n+1}}{(2az - 1)^{n+1}}, \end{aligned}$$

$$\begin{aligned}
y_{20n-9} &= \frac{(-1)^n q(2eq-1)^n}{(eq-1)^{2n}}, & y_{20n-8} &= \frac{(-1)^n p(2dp-1)^n}{(dp-1)^{2n}}, & y_{20n-7} &= \frac{(-1)^n o(2co-1)^n}{(co-1)^{2n}}, \\
y_{20n-6} &= \frac{(-1)^n L(2Lb-1)^n}{(Lb-1)^{2n}}, & y_{20n-5} &= \frac{(-1)^n z(2az-1)^n}{(az-1)^{2n}}, & y_{20n-4} &= (-1)^n t(s^2t^2-1)^n, \\
y_{20n-3} &= (-1)^n w(k^2w^2-1)^n, & y_{20n-2} &= (-1)^n v(h^2v^2-1)^n, & y_{20n-1} &= (-1)^n u(g^2u^2-1)^n, \\
y_{20n} &= (-1)^n r(f^2r^2-1)^n, & y_{20n+1} &= \frac{(-1)^n q(2eq-1)^n}{(eq-1)^{2n+1}}, & y_{20n+2} &= \frac{(-1)^n p(2dp-1)^n}{(dp-1)^{2n+1}}, \\
y_{20n+3} &= \frac{(-1)^n o(2co-1)^n}{(co-1)^{2n+1}}, & y_{20n+4} &= \frac{(-1)^n L(2Lb-1)^n}{(Lb-1)^{2n+1}}, & y_{20n+5} &= \frac{(-1)^n z(2az-1)^n}{(az-1)^{2n+1}}, \\
y_{20n+6} &= (-1)^{n+1} t(st+1)(s^2t^2-1)^n, & y_{20n+7} &= (-1)^{n+1} w(kw+1)(k^2w^2-1)^n, \\
y_{20n+8} &= (-1)^{n+1} v(hv+1)(h^2v^2-1)^n, & y_{20n+9} &= (-1)^{n+1} u(gu+1)(g^2u^2-1)^n, \\
y_{20n+10} &= (-1)^{n+1} r(fr+1)(f^2r^2-1)^n.
\end{aligned}$$

Proof. As the proof of Theorem 1 and will be left to the reader.

Theorem 4. Consider $\{x_n, y_n\}$ are solutions of the system (4.2) with $x_{-9}y_{-4}, x_{-8}y_{-3}, x_{-7}y_{-2}, x_{-6}y_{-1}, x_{-5}y_0 \neq -1, \neq -\frac{1}{2}$ and $x_{-4}y_{-9}, x_{-3}y_{-8}, x_{-2}y_{-7}, x_{-1}y_{-6}, x_0y_{-5} \neq \pm 1$. Then for $n = 0, 1, 2, \dots$,

$$\begin{aligned}
x_{20n-9} &= \frac{s(2st+1)^n}{(st+1)^{2n}}, & x_{20n-8} &= \frac{k(2kw+1)^n}{(kw+1)^{2n}}, & x_{20n-7} &= \frac{h(2hv+1)^n}{(hv+1)^{2n}}, \\
x_{20n-6} &= \frac{g(2gu+1)^n}{(gu+1)^{2n}}, & x_{20n-5} &= \frac{f(2fr+1)^n}{(fr+1)^{2n}}, & x_{20n-4} &= (-1)^n e(e^2q^2-1)^n, \\
x_{20n-3} &= (-1)^n d(d^2p^2-1)^n, & x_{20n-2} &= (-1)^n c(c^2o^2-1)^n, & x_{20n-1} &= (-1)^n b(L^2b^2-1)^n, \\
x_{20n} &= (-1)^n a(a^2z^2-1)^n, & x_{20n+1} &= \frac{s(2st+1)^n}{(st+1)^{2n+1}}, & x_{20n+2} &= \frac{k(2kw+1)^n}{(kw+1)^{2n+1}}, \\
x_{20n+3} &= \frac{h(2hv+1)^n}{(hv+1)^{2n+1}}, & x_{20n+4} &= \frac{g(2gu+1)^n}{(gu+1)^{2n+1}}, & x_{20n+5} &= \frac{f(2fr+1)^n}{(fr+1)^{2n+1}}, \\
x_{20n+6} &= (-1)^n e(eq+1)(e^2q^2-1)^n, & x_{20n+7} &= (-1)^n d(dp+1)(d^2p^2-1)^n, \\
x_{20n+8} &= (-1)^n c(co+1)(c^2o^2-1)^n, & x_{20n+9} &= (-1)^n b(Lb+1)(L^2b^2-1)^n, \\
x_{20n+10} &= (-1)^n a(az+1)(a^2z^2-1)^n,
\end{aligned}$$


$$\begin{aligned}
y_{20n-9} &= \frac{(-1)^n q}{(e^2q^2-1)^n}, & y_{20n-8} &= \frac{(-1)^n p}{(d^2p^2-1)^n}, & y_{20n-7} &= \frac{(-1)^n o}{(c^2o^2-1)^n}, \\
y_{20n-6} &= \frac{(-1)^n L}{(L^2b^2-1)^n}, & y_{20n-5} &= \frac{(-1)^n z}{(a^2z^2-1)^n}, & y_{20n-4} &= \frac{t(st+1)^{2n}}{(2st+1)^n}, \\
y_{20n-3} &= \frac{w(kw+1)^{2n}}{(2kw+1)^n}, & y_{20n-2} &= \frac{v(hv+1)^{2n}}{(2hv+1)^n}, & y_{20n-1} &= \frac{u(gu+1)^{2n}}{(2gu+1)^n}, \\
y_{20n} &= \frac{r(fr+1)^{2n}}{(2fr+1)^n}, & y_{20n+1} &= \frac{(-1)^n q(2eq-1)^n}{(eq-1)^{2n+1}}, & y_{20n+2} &= \frac{(-1)^n p(2dp-1)^n}{(dp-1)^{2n+1}}, \\
y_{20n+3} &= \frac{(-1)^n o(2co-1)^n}{(co-1)^{2n+1}}, & y_{20n+4} &= \frac{(-1)^n L(2Lb-1)^n}{(Lb-1)^{2n+1}}, & y_{20n+5} &= \frac{(-1)^n z(2az-1)^n}{(az-1)^{2n+1}}, \\
y_{20n+6} &= \frac{-t(st+1)^{2n+1}}{(2st+1)^{n+1}}, & y_{20n+7} &= \frac{-w(kw+1)^{2n+1}}{(2kw+1)^{n+1}}, & y_{20n+8} &= \frac{-v(hv+1)^{2n+1}}{(2hv+1)^{n+1}}, \\
y_{20n+9} &= \frac{-u(gu+1)^{2n+1}}{(2gu+1)^{n+1}}, & y_{20n+10} &= \frac{-r(fr+1)^{2n+1}}{(2fr+1)^{n+1}}.
\end{aligned}$$

Proof. As in the proof of Theorem 2 and will be omitted.


4.2. Numerical Examples

In this subsection to support our theoretical discussions, we consider some interesting numerical examples.


Example 7. When we put the initial conditions $x_{-9} = -2.1$, $x_{-8} = 0.3$, $x_{-7} = -4$, $x_{-6} = 0.11$, $x_{-5} = 0.5$, $x_{-4} = 1.9$, $x_{-3} = -1$, $x_{-2} = 0.039$, $x_{-1} = 0.2$, $x_0 = -0.7$, $y_{-9} = 0.35$, $y_{-8} = -0.38$, $y_{-7} = 0.9$, $y_{-6} = 0.1$, $y_{-5} = -0.14$, $y_{-4} = -0.11$, $y_{-3} = 0.6$, $y_{-2} = -0.5$, $y_{-1} = 0.1$, $y_0 = 0.62$ for system (4). (See Fig. 7).



Example 8. See Figure 8 it shows the periodicity of the solutions for system (4) where the initial conditions takes the numbers $x_{-9} = x_{-8} = x_{-7} = x_{-6} = x_{-5} = 0$, $x_{-4} = 1.9$, $x_{-3} = -1$, $x_{-2} = 0.39$, $x_{-1} = 0.2$, $x_0 = 0.7$, $y_{-9} = y_{-8} = y_{-7} = y_{-6} = y_{-5} = 0$, $y_{-4} = 0.11$, $y_{-3} = 0.6$, $y_{-2} = 0.5$, $y_{-1} = 0.81$, $y_0 = 0.62$.



Example 9. Let we take for the difference system (5) the initial conditions $x_{-9} = 0.8$, $x_{-8} = -0.2$, $x_{-7} = 0.25$, $x_{-6} = -0.33$, $x_{-5} = 0.15$, $x_{-4} = 0.9$, $x_{-3} = -0.01$, $x_{-2} = 0.39$, $x_{-1} = 0.2$, $x_0 = 0.7$, $y_{-9} = 0.085$, $y_{-8} = 0.3$, $y_{-7} = -0.29$, $y_{-6} = 0.56$, $y_{-5} = 0.41$, $y_{-4} = -0.11$, $y_{-3} = 0.6$, $y_{-2} = 0.5$, $y_{-1} = 0.81$, $y_0 = 0.62$. (See Fig. 9).



References

1. Asiri, A., El-Dessoky, M. M. and Elsayed, E. M., *Solution of a third order fractional system of difference equations*, Journal of Computational Analysis and Applications, 24 (3),444-453 (2018).
2. Alghamdi, M., Elsayed, E. M. and El-Dessoky, M. M., *On the Solutions of Some Systems of Second Order Rational Difference Equations*, Life Science Journal, 10 (3), 344-351 (2013).

3. Clark, D. and Kulenovic, M. R. S., *A coupled system of rational difference equations*, Computers & Mathematics with Applications, 43, 849-867 (2002).
4. Din, Q. and Elsayed, E. M., *Stability analysis of a discrete ecological model*, Computational Ecology and Software, 4 (2) , 89-103(2014).
5. Elabbasy, E. M., Elsadany, A. A. and Zhang, Y., *Bifurcation analysis and chaos in a discrete reduced Lorenz system*, Applied Mathematics and Computation, 228, 184-194 (2014).
6. Elabbasy, E. M., Elsadany, A. A. and Ibrahim, S., *Behavior and periodic solutions of a two-dimensional systems of rational difference equations*, Journal of Interpolation and Approximation in Scientific Computing, 2016 (2),, 18 pages (2016).
7. El-Dessoky, M. M., *The form of solutions and periodicity for some systems of third - order rational difference equations*, Mathematical Methods in the Applied Sciences 39 (5), 1076-1092, (2016).
8. El-Dessoky, M. M., *On a solvable for some systems of rational difference equations*, Journal of Computational Analysis and Applications 9 (6), 3744-3759, (2016).
9. El-Dessoky, M. M., *On a Solvable for Some Nonlinear Systems of Difference Equations*, Journal of Computational and Theoretical Nanoscience 12 (11), 3432-3442, (2015).
10. El-Dessoky, M. M. and Elsayed, E. M., *On a solution of system of three fractional difference equations*, Journal of Computational Analysis and Applications 19 (4), 760-769, (2015).
11. El-Metwally, H. and Elsayed, E. M., *Form of solutions and periodicity for systems of difference equations*, Journal of Computational Analysis and Applications 15 (5), 852-857, (2013).
12. Elsayed, E. M., *Solutions of Rational Difference System of Order Two*, Mathematical and Computer Modelling 55, 378—384, (2012).
13. Elsayed, E. M., *On the solutions and periodic nature of some systems of difference equations*, International Journal of Biomathematics 7 (6), 1450067, (26 pages), (2014).
14. Elsayed, E. M., *Solution for systems of difference equations of rational form of order two*, Computational and Applied Mathematics 33 (3), 751-765, (2014).
15. Elsayed, E. M., Alotaibi, A. and Almaylubi, H. A., *On a solutions of fourth order rational systems of difference equations*, Journal of Computational Analysis and applications 22 (7), 1298-1308, (2017).
16. Elsayed, E. M. and Ahmed, A. M., *Dynamics of a three-dimensional systems of rational difference equations*, Mathematical Methods in the Applied Sciences 39 (5), 1026-1038, (2016).
17. Elsayed, E. M., El-Dessoky, M. M. and Alotaibi, A., *On the Solutions of a General System of Difference Equations*, Discrete Dynamics in Nature and Society Vol. 2012, Article ID 892571, 12 pages, (2012).
18. Halim, Y., Touafek, N. and Elsayed, E. M., *Closed form solutions of some systems of rational difference equations in terms of Fibonacci numbers*, Dynamics of Continuous, Discrete and Impulsive Systems, Series A: Mathematical Analysis 21, 473-486, (2014).
19. Ibrahim, T. F. and Touafek, N., *Max-type system of difference equations with positive two-periodic sequences*, Mathematical Methods in the Applied Sciences 37, 2541-2553, (2014).
20. Khan, A. Q., Qureshi, M. N. and Din, Q., *Qualitative behavior of two systems of second-order rational difference equations*, Journal of Computational Analysis and Applications 18 (6), 1027-1041, (2015).
21. Kurbanli, A. S., Cinar, C. and Erdoan, M., *On the periodicity of solutions of the system of rational difference equations*, Applied Mathematics 2 (4), 410-413, (2011).
22. Kurbanli, A. S., Cinar, C. and Yalçınkaya, I., *On the behavior of positive solutions of the system of rational difference equations*, Mathematical and Computer Modelling 53, 1261-1267, (2011).
23. Ma, H. and Feng, H., *On Positive Solutions for the Rational Difference Equation Systems*, International Scholarly Research Notices Volume 2014, Article ID 857480, 4 pages, (2014).
24. Mansour, M., El-Dessoky, M. M. and Elsayed, E. M., *On the solution of rational systems of difference equations*, Journal of Computational Analysis and Applications 15 (5), 967-976, (2013).
25. Tollu, D., Yazlik, Y. and Taskara, N., *The Solutions of Four Riccati Difference Equations Associated with Fibonacci Numbers*, Balkan Journal of Mathematics 2, 163-172, (2014).
26. Touafek, N. and Elsayed, E. M., *On the solutions of systems of rational difference equations*, Mathematical and Computer Modelling 55, 1987-1997, (2012).
27. Touafek, N. and Elsayed, E. M., *On the periodicity of some systems of nonlinear difference equations*, Bulletin Mathématique de la Société des Sciences Mathématiques de Roumanie 55 (103) (2), 217-224, (2012).
28. Touafek, N. and Elsayed, E. M., *On a second order rational systems of difference equation*, Hokkaido Mathematical Journal 44 (1), 29-45, (2015).

29. Yazlik, Y., Tollu, D. T. and Taskara, N., *On the Behaviour of Solutions for Some Systems of Difference Equations*, Journal of Computational Analysis and Applications 18 (1), 166-178, (2015).
30. Zhang, Q., Liu, J. and Luo, Z., *Dynamical Behavior of a System of Third-Order Rational Difference Equation*, Discrete Dynamics in Nature and Society Volume 2015, Article ID 530453, 6 pages, (2015).

E. M. Elsayed,
Department of Mathematics,
King AbdulAziz University, Faculty of Science,
P. O. Box 80203, Jeddah 21589, Saudi Arabia.

and

E. M. Elsayed,
Department of Mathematics,
Faculty of Science, Mansoura University,
Mansoura 35516, Egypt.

and

H. S. Gafel,
Department of Mathematics,
King AbdulAziz University, Faculty of Science,
P. O. Box 80203, Jeddah 21589, Saudi Arabia.
E-mail address: emelsayed@mans.edu.eg, emmelsayed@yahoo.com

and

H. S. Gafel,
Department of Mathematics and Statistics, College of Science,
Taif University, P.O. Box 11099, Taif 21944, Saudi Arabia.
E-mail address: emmelsayed@yahoo.com, h.gafal@tu.edu.sa