



On an elliptic equation of p -Laplacian type with nonlinear boundary condition

Nguyen Thanh Chung

ABSTRACT: We consider elliptic equations of p -Laplacian type with the nonlinear boundary condition of the form

$$\begin{cases} -\Delta_p u + |u|^{p-2}u &= \lambda f_1(u) + \mu g_1(u) & \text{in } \Omega, \\ |\nabla u|^{p-2} \frac{\partial u}{\partial n} &= \lambda f_2(u) + \mu g_2(u) & \text{in } \partial\Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^N$ ($N \geq 3$) is a bounded domain with smooth boundary $\partial\Omega$, $\frac{\partial}{\partial n}$ is the outer unit normal derivative, λ, μ are parameters. The functions f_i , $i = 1, 2$, are assumed to be $(p-1)$ -sublinear while g_i , $i = 1, 2$, are $(p-1)$ -asymptotically linear at infinity. Using variational techniques, an existence result is given.

Key Words: Elliptic equation, p -Laplacian type, $(p-1)$ -sublinear, $(p-1)$ -asymptotically linear, Nonlinear boundary condition.

Contents

1 Introduction	43
2 Existence of solutions	45

1. Introduction

Consider the elliptic equation of p -Laplacian type with nonlinear boundary condition

$$\begin{cases} -\Delta_p u + |u|^{p-2}u &= \lambda f_1(u) + \mu g_1(u) & \text{in } \Omega, \\ |\nabla u|^{p-2} \frac{\partial u}{\partial n} &= \lambda f_2(u) + \mu g_2(u) & \text{in } \partial\Omega, \end{cases} \quad (1.1)$$

where $\Omega \subset \mathbb{R}^N$ ($N \geq 3$) is a bounded domain with smooth boundary $\partial\Omega$, $\frac{\partial}{\partial n}$ is the outer unit normal derivative, $1 < p < N$, λ, μ are parameters.

Problem (1.1) has been studied in many works, such as [1,2,3,4,5,9], in which the authors have used different methods to obtain the existence of solutions. In a recent paper [7], we have considered the situation: $g_i \equiv 0$ ($i = 1, 2$), f_i , $i = 1, 2$, are $(p-1)$ -sublinear at infinity. We then used the three critical point theorem of G. Bonanno [6] to obtain a multiplicity result for (1.1). A natural question is to see what happens if the problem in [7] is affected by a certain perturbation. For this purpose, in this note, we establish an existence result for (1.1) in the case when $f_i : \mathbb{R} \rightarrow \mathbb{R}$, $i = 1, 2$, are $(p-1)$ -sublinear and $g_i : \mathbb{R} \rightarrow \mathbb{R}$, $i = 1, 2$, are

2000 *Mathematics Subject Classification*: 35J65, 35J20.

$(p-1)$ -asymptotically at infinity. The proof relies essentially on the minimum principle in [8, Theorem 2.1].

In order to state the main result of this work, we would introduce the following hypotheses

(f) $f_i, i = 1, 2$ are continuous and $(p-1)$ -sublinear at infinity, i.e.,

$$\lim_{|t| \rightarrow \infty} \frac{|f_i(t)|}{|t|^{p-1}} = 0;$$

(g) $g_i, i = 1, 2$ are continuous and $(p-1)$ -asymptotically at infinity, i.e.,

$$\lim_{|t| \rightarrow \infty} \frac{|g_i(t)|}{|t|^{p-1}} = l_i < +\infty.$$

Let $W^{1,p}(\Omega)$ be the usual Sobolev space with respect to the norm

$$\|u\|_{1,p}^p = \int_{\Omega} (|\nabla u|^p + |u|^p) dx$$

and $W_0^{1,p}(\Omega)$ is the closure of $C_0^\infty(\Omega)$ in $W^{1,p}(\Omega)$. For any $1 < p < N$ and $1 \leq q \leq p^* = \frac{Np}{N-p}$, we denote by $S_{q,\Omega}$ the best constant in the embedding $W^{1,p}(\Omega) \hookrightarrow L^q(\Omega)$ and for all $1 \leq q \leq p_* = \frac{(N-1)p}{N-p}$, we also denote by $S_{q,\partial\Omega}$ the best constant in the embedding $W^{1,p}(\Omega) \hookrightarrow L^q(\partial\Omega)$, i.e.

$$S_{q,\partial\Omega} = \inf_{u \in W^{1,p}(\Omega) \setminus W_0^{1,p}(\Omega)} \frac{\int_{\Omega} (|\nabla u|^p + |u|^p) dx}{\left(\int_{\partial\Omega} |u|^q d\sigma \right)^{\frac{p}{q}}}.$$

Moreover, if $1 \leq q < p^*$, then the embedding $W^{1,p}(\Omega) \hookrightarrow L^q(\Omega)$ is compact and if $1 \leq q < p_*$, then the embedding $W^{1,p}(\Omega) \hookrightarrow L^q(\partial\Omega)$ is compact. As a consequence, we have the existence of extremals, i.e. functions where the infimum is attained (see [2,5]).

Definition 1.1. A function $u \in W^{1,p}(\Omega)$ is said to be a weak solution of problem (1.1) if and only if

$$\begin{aligned} \int_{\Omega} \left[|\nabla u|^{p-2} \nabla u \cdot \nabla v + |u|^{p-2} uv \right] dx - \lambda \int_{\Omega} f_1(u) v dx - \lambda \int_{\partial\Omega} f_2(u) v d\sigma \\ - \mu \int_{\Omega} g_1(u) v dx - \mu \int_{\partial\Omega} g_2(u) v d\sigma = 0 \end{aligned}$$

for all $v \in W^{1,p}(\Omega)$.

Theorem 1.2. Assume conditions (f) and (g) are fulfilled. Moreover, there exists $s_0 > 0$ such that

$$F_1(s_0) := \int_0^{s_0} f_1(t) dt > 0 \text{ and } F_2(s_0) := \int_0^{s_0} f_2(t) dt > 0.$$

Then for each $\lambda \in \mathbb{R}$ large enough, there exists $\bar{\mu} > 0$, such that problem (1.1) has at least a non-trivial weak solution u in $W^{1,p}(\Omega)$ for every $\mu \in (0, \bar{\mu})$.

2. Existence of solutions

For $\lambda, \mu \in \mathbb{R}$, let us define the functional $J_{\lambda, \mu} : W^{1,p}(\Omega) \rightarrow \mathbb{R}$ associated to problem (1.1) by the formula

$$\begin{aligned} J_{\lambda, \mu}(u) &= \frac{1}{p} \int_{\Omega} [|\nabla u|^p + |u|^p] dx - \lambda \int_{\Omega} F_1(u) dx - \lambda \int_{\partial\Omega} F_2(u) d\sigma \\ &\quad - \mu \int_{\Omega} G_1(u) dx - \mu \int_{\partial\Omega} G_2(u) d\sigma \\ &= \Lambda(u) - I_{\lambda, \mu}(u), \end{aligned} \quad (2.1)$$

where

$$\begin{aligned} \Lambda(u) &= \frac{1}{p} \int_{\Omega} [|\nabla u|^p + |u|^p] dx, \\ I_{\lambda, \mu}(u) &= \lambda \int_{\Omega} F_1(u) dx + \lambda \int_{\partial\Omega} F_2(u) d\sigma + \mu \int_{\Omega} G_1(u) dx + \mu \int_{\partial\Omega} G_2(u) d\sigma \end{aligned} \quad (2.2)$$

for all $u \in W^{1,p}(\Omega)$. Then, a simple computation shows that $J_{\lambda, \mu}$ is of C^1 class and

$$\begin{aligned} DJ_{\lambda, \mu}(u)(v) &= \int_{\Omega} [|\nabla u|^{p-2} \nabla u \cdot \nabla v + |u|^{p-2} uv] dx - \lambda \int_{\Omega} f_1(u) v dx - \lambda \int_{\partial\Omega} f_2(u) v d\sigma \\ &\quad - \mu \int_{\Omega} g_1(u) v dx - \mu \int_{\partial\Omega} g_2(u) v d\sigma = 0 \end{aligned}$$

for all $u, v \in W^{1,p}(\Omega)$. Thus, weak solutions of problem (1.1) are exactly the critical points of $J_{\lambda, \mu}$.

Lemma 2.1. *For every $\lambda \in \mathbb{R}$, there exists $\bar{\mu} > 0$, depending on λ , such that for every $\mu \in (0, \bar{\mu})$, the functional $J_{\lambda, \mu}$ is coercive.*

Proof. Firstly, we have

$$S_{p, \Omega} \|u\|_{L^p(\Omega)} \leq \|u\|_{1,p} \text{ and } S_{p, \partial\Omega} \|u\|_{L^p(\partial\Omega)} \leq \|u\|_{1,p}$$

for all $u \in W^{1,p}(\Omega)$.

Let us fix $\lambda \in \mathbb{R}$, arbitrary. By (f), there exist $\delta_i = \delta_i(\lambda)$, $i = 1, 2$, such that

$$|f_1(t)| \leq S_{p, \Omega}^p \frac{1}{2(1+|\lambda|)} |t|^{p-1}, \quad \forall |t| \geq \delta_1$$

and

$$|f_2(t)| \leq S_{p, \partial\Omega}^p \frac{1}{2(1+|\lambda|)} |t|^{p-1}, \quad \forall |t| \geq \delta_2.$$

Integrating the above inequalities, we have

$$|F_1(t)| \leq S_{p, \Omega}^p \frac{1}{2p(1+|\lambda|)} |t|^p + \max_{|s| \leq \delta_1} |f_1(s)| |t|, \quad \forall t \in \mathbb{R} \quad (2.3)$$

and

$$|F_2(t)| \leq S_{p,\partial\Omega}^p \frac{1}{2p(1+|\lambda|)} |t|^p + \max_{|s| \leq \delta_2} |f_2(s)| |t|, \quad \forall t \in \mathbb{R}. \quad (2.4)$$

Since g_i , $i = 1, 2$ are $(p-1)$ -asymptotically linear at infinity, there exist two constants $m_i > 0$, $i = 1, 2$, such that

$$|g_1(t)| \leq m_1 p S_{p,\Omega}^p |t|^{p-1} + m_1,$$

$$|g_2(t)| \leq m_2 p S_{p,\partial\Omega}^p |t|^{p-1} + m_2$$

for all $t \in \mathbb{R}$. It implies that

$$|G_1(t)| \leq m_1 S_{p,\Omega}^p |t|^p + m_1 |t|, \quad (2.5)$$

and

$$|G_2(t)| \leq m_2 S_{p,\partial\Omega}^p |t|^p + m_2 |t| \quad (2.6)$$

for all $t \in \mathbb{R}$.

Hence, for any $u \in W^{1,p}(\Omega)$, we deduce that

$$\begin{aligned} J_{\lambda,\mu}(u) &\geq \Lambda(u) - |I_{\lambda,\mu}(u)| \\ &\geq \frac{1}{p} \|u\|_{1,p}^p - \frac{|\lambda|}{2p(1+|\lambda|)} \|u\|_{1,p}^p - \frac{|\lambda|}{S_{p,\Omega}} |\Omega|_{N'}^{\frac{1}{p'}} \|u\|_{1,p} \max_{|s| \leq \delta_1} |f_1(s)| \\ &\quad - \frac{|\lambda|}{2p(1+|\lambda|)} \|u\|_{1,p}^p - \frac{|\lambda|}{S_{p,\partial\Omega}} |\partial\Omega|_{N-1}^{\frac{1}{p'}} \|u\|_{1,p} \max_{|s| \leq \delta_1} |f_2(s)| \\ &\quad - |\mu| m_1 \|u\|_{1,p}^p - m_1 \frac{|\mu|}{S_{p,\Omega}} |\Omega|_{N'}^{\frac{1}{p'}} \|u\|_{1,p} \\ &\quad - |\mu| m_2 \|u\|_{1,p}^p - m_2 \frac{|\mu|}{S_{p,\partial\Omega}} |\partial\Omega|_{N-1}^{\frac{1}{p'}} \|u\|_{1,p} \\ &= \left(\frac{1}{p(1+|\lambda|)} - |\mu|(m_1 + m_2) \right) \|u\|_{1,p}^p - \frac{|\lambda|}{S_{p,\Omega}} |\Omega|_{N'}^{\frac{1}{p'}} \|u\|_{1,p} \max_{|s| \leq \delta_1} |f_1(s)| \\ &\quad - \frac{|\lambda|}{S_{p,\partial\Omega}} |\partial\Omega|_{N-1}^{\frac{1}{p'}} \|u\|_{1,p} \max_{|s| \leq \delta_1} |f_2(s)| - m_1 \frac{|\mu|}{S_{p,\Omega}} |\Omega|_{N'}^{\frac{1}{p'}} \|u\|_{1,p} \\ &\quad - m_2 \frac{|\mu|}{S_{p,\partial\Omega}} |\partial\Omega|_{N-1}^{\frac{1}{p'}} \|u\|_{1,p}, \end{aligned}$$

where $p' = \frac{p}{p-1}$. Let $\bar{\mu} = \frac{1}{p(m_1+m_2)(1+|\lambda|)}$ and fix $\mu \in (0, \bar{\mu})$. Since $p > 1$ we have $J_{\lambda,\mu}(u) \rightarrow +\infty$ as $\|u\|_{1,p} \rightarrow \infty$. Thus, the functional $J_{\lambda,\mu}$ is coercive. \square

Lemma 2.2. *Let λ and $\bar{\mu}$ be chosen as in the previous lemma. Then for each $\mu \in (0, \bar{\mu})$, the functional $J_{\lambda,\mu}$ satisfies the Palais-Smale condition.*

Proof. Let $\{u_m\}$ be a sequence in $W^{1,p}(\Omega)$ such that

$$J_{\lambda,\mu}(u_m) \rightarrow \bar{c}, \quad DJ_{\lambda,\mu}(u_m) \rightarrow 0 \text{ in } W^{-1,p}(\Omega) \text{ as } m \rightarrow \infty. \quad (2.7)$$

Since the functional $J_{\lambda,\mu}$ is coercive, the sequence $\{u_m\}$ is bounded in $W^{1,p}(\Omega)$. Then, there exist a subsequence still denoted by $\{u_m\}$ and a function $u \in W^{1,p}(\Omega)$, such that $\{u_m\}$ converges weakly to u in $W^{1,p}(\Omega)$. Hence, $\{\|u_m - u\|_{1,p}\}$ is bounded and by (2.7), $DJ_{\lambda,\mu}(u_m)(u_m - u)$ converges to 0 as $m \rightarrow \infty$.

By (f), there exists a constant $C_1 > 0$ such that

$$|f_i(t)| \leq C_1(1 + |t|^{p-1}), \quad i = 1, 2$$

for all $t \in \mathbb{R}$. Therefore,

$$\begin{aligned} 0 &\leq \int_{\Omega} |f_1(u_m)| |u_m - u| dx \leq C_1 \int_{\Omega} |u_m - u| dx + C \int_{\Omega} |u_m|^{p-1} |u_m - u| dx \\ &\leq C_1 \left[|\Omega|^{\frac{1}{p'}} + \|u_m\|_{L^p(\Omega)}^{p-1} \right] \|u_m - u\|_{L^p(\Omega)} \end{aligned}$$

and

$$\begin{aligned} 0 &\leq \int_{\partial\Omega} |f_2(u_m)| |u_m - u| dx \leq C_1 \int_{\partial\Omega} |u_m - u| dx + C_1 \int_{\partial\Omega} |u_m|^{p-1} |u_m - u| dx \\ &\leq C_1 \left[|\partial\Omega|^{\frac{1}{p'}} + \|u_m\|_{L^p(\partial\Omega)}^{p-1} \right] \|u_m - u\|_{L^p(\partial\Omega)}. \end{aligned}$$

Since $\{u_m\}$ converges strongly to u in the spaces $L^p(\Omega)$ and $L^p(\partial\Omega)$, the above inequalities imply that

$$\lim_{m \rightarrow \infty} \int_{\Omega} f_1(u_m)(u_m - u) dx = 0 \quad (2.8)$$

and

$$\lim_{m \rightarrow \infty} \int_{\partial\Omega} f_2(u_m)(u_m - u) dx = 0. \quad (2.9)$$

On the other hand, by (g), there exists a constant $C_2 > 0$ such that

$$|g_i(t)| \leq C_2(1 + |t|^{p-1}), \quad i = 1, 2$$

for all $t \in \mathbb{R}$. Therefore, the similar arguments above show that

$$\lim_{m \rightarrow \infty} \int_{\Omega} g_1(u_m)(u_m - u) dx = 0 \quad (2.10)$$

and

$$\lim_{m \rightarrow \infty} \int_{\partial\Omega} g_2(u_m)(u_m - u) dx = 0. \quad (2.11)$$

By relations (2.8)-(2.11), we get

$$\lim_{m \rightarrow \infty} DI_{\lambda,\mu}(u_m)(u_m - u) = 0. \quad (2.12)$$

Combining (2.11) and (2.7), it follows that

$$\lim_{m \rightarrow \infty} \Lambda(u_m)(u_m - u) = 0. \quad (2.13)$$

Hence, standard arguments help us to show that the sequence $\{u_m\}$ converges strongly to u in $W^{1,p}(\Omega)$. Thus, the functional $J_{\lambda,\mu}$ satisfies the Palais-Smale condition in $W^{1,p}(\Omega)$. \square

Proof Theorem 1.2. By Lemmas 2.1 and 2.2, using the minimum principle [8, Theorem 2.1], we deduce that for each $\lambda \in \mathbb{R}$, there exists $\bar{\mu} > 0$, such that for any $\mu \in (0, \bar{\mu})$, problem (1.1) has a weak solution $u \in W^{1,p}(\Omega)$. We will show that u is not trivial for λ large enough. Indeed, let s_0 be a real number such that

$$F_1(s_0) := \int_0^{s_0} f_1(t)dt > 0 \text{ and } F_2(s_0) := \int_0^{s_0} f_2(t)dt > 0$$

and let $\Omega_0 \subset \Omega$ be an open subset with $|\Omega_0|_N > 0$. Then, there exists $u_0 \in C_0^\infty(\Omega)$ such that $u_0(x) \equiv s_0$ on Ω_0 and $0 \leq u_0(x) \leq s_0$ in $\Omega \setminus \Omega_0$. We have

$$\begin{aligned} J_{\lambda,\mu}(u_0) &= \frac{1}{p} \int_{\Omega} [|\nabla u_0|^p + |u_0|^p] dx - \lambda \int_{\Omega} F_1(u_0) dx - \lambda \int_{\partial\Omega} F_2(u_0) dx \\ &\quad - \mu \int_{\Omega} G_1(u_0) dx - \mu \int_{\partial\Omega} G_2(u_0) dx \\ &\leq \frac{1}{p} \int_{\Omega} [|\nabla u_0|^p + |u_0|^p] dx - \lambda \int_{\Omega_0} F_1(u_0) dx - \lambda \int_{\partial\Omega_0} F_2(u_0) dx \\ &\quad - \mu \int_{\Omega_0} G_1(u_0) dx - \mu \int_{\partial\Omega_0} G_2(u_0) dx \\ &= C - \lambda \left(F_1(s_0) |\Omega_0|_N + F_2(s_0) |\Omega_0|_{N-1} \right), \end{aligned}$$

where C is a positive constant (C depends on μ). Therefore, for $\lambda > 0$ large enough, we have $J_{\lambda,\mu}(u_0) < 0$. Thus, the solution u is not trivial. The proof of Theorem 1.2 is now completed. \square

References

1. E. A. M. ABREU, J. M. DO Ó AND E.S. MEDEIROS, Multiplicity of solutions for a class of quasilinear nonhomogeneous Neumann problems, *Nonlinear Anal.*, **60** (2005), 1443-1471.
2. J.F. BONDER, Multiple positive solutions for quasilinear elliptic problems with sign-changing nonlinearities, *Abstr. Appl. Anal.*, **2004**(12) (2004), 1047-1056.
3. J.F. BONDER, Multiple solutions for the p -Laplacian equation with nonlinear boundary conditions, *Electron. J. Diff. Equ.*, **2006**(37) (2006), 1-7.
4. J.F. BONDER, S. MARTÍNEZ AND J.D. ROSSI, Existence results for gradient elliptic systems with nonlinear boundary conditions, *Nonlinear Differ. Eqns. Appl. (NoDEA)*, **14**(1-2) (2007), 153-179.
5. J.F. BONDER AND J.D. ROSSI, Existence results for the p -Laplacian with nonlinear boundary conditions, *J. Math. Anal. Appl.*, **263** (2001), 195-223.

6. G. BONANNO, Some remarks on a three critical points theorem, *Nonlinear Anal.*, **54** (2003), 651-665.
7. N.T. CHUNG, Multiple solutions for quasilinear elliptic problems with nonlinear boundary conditions, *Electron. J. Diff. Equ.*, **2008**(165) (2008), 1-6.
8. A. ANANE, O. CHAKRONE, B. KARIM AND A. ZEROUALI, Existence of solutions for a resonant Steklov Problem, *Bol. Soc. Paranaense de Mat.*, **27**(1) (2009), 85-90.
9. J.H. ZHAO AND P.H. ZHAO, Infinitely many weak solutions for a p -Laplacian equations with nonlinear boundary conditions, *Electron. J. Diff. Equ.*, **2006** (90) (2006), 1-14.

Nguyen Thanh Chung,
Department of Mathematics and Informatics,
Quang Binh University, 312 Ly Thuong Kiet,
Dong Hoi, Vietnam
E-mail address: chungnt.qbu@gmail.com