



Biharmonic Curves according to Parallel Transport Frame in \mathbb{E}^4

Talat Körpınar and Essin Turhan

ABSTRACT: In this paper, we study biharmonic curves according to parallel transport frame in \mathbb{E}^4 . We give some characterizations for curvatures of a biharmonic curve in \mathbb{E}^4 .

Key Words: Heisenberg group, fluid flow, biharmonic curve, parallel transport frame.

Contents

1 Introduction	213
2 Preliminaries	213
3 Biharmonic curves according to parallel transport frame in \mathbb{E}^4	215

1. Introduction

In the last decade there has been a growing interest in the theory of biharmonic maps which can be divided in two main research directions. On the one side, constructing the examples and classification results have become important from the differential geometric aspect. The other side is the analytic aspect from the point of view of partial differential equations, because biharmonic maps are solutions of a fourth order strongly elliptic semilinear PDE.

In this paper, we study biharmonic curves according to parallel transport frame in \mathbb{E}^4 . We give some characterizations for curvatures of a biharmonic curve in \mathbb{E}^4 .

2. Preliminaries

To meet the requirements in the next sections, here, the basic elements of the theory of curves in the space \mathbb{E}^4 are briefly presented.

The Euclidean 4-space \mathbb{E}^4 provided with the standard flat metric given by

$$\langle, \rangle = dx_1^2 + dx_2^2 + dx_3^2 + dx_4^2,$$

where (x_1, x_2, x_3, x_4) is a rectangular coordinate system of \mathbb{E}^4 . Recall that, the norm of an arbitrary vector $a \in \mathbb{E}^4$ is given by $\|a\| = \sqrt{\langle a, a \rangle}$. γ is called a unit speed curve if velocity vector v of γ satisfies $\|v\| = 1$.

Let $\{\mathbf{T}, \mathbf{N}, \mathbf{B}_1, \mathbf{B}_2\}$ denotes the moving Frenet frame of the unit speed curve γ , where $\mathbf{T}, \mathbf{N}, \mathbf{B}_1$ and \mathbf{B}_2 are the tangent, the principal normal, the first binormal and the second binormal vector fields, respectively. Then the Serret-Frenet formulae

2000 *Mathematics Subject Classification*: 31B30, 58E20

are given by binormal and the second binormal vector fields, respectively. Then the Serret-Frenet formulae are given by

$$\begin{aligned}\mathbf{T}' &= \kappa\mathbf{N} \\ \mathbf{N}' &= -\kappa\mathbf{T} + \tau\mathbf{B}_1 \\ \mathbf{B}'_1 &= -\tau\mathbf{N} + \sigma\mathbf{B}_2 \\ \mathbf{B}'_2 &= -\sigma\mathbf{B}_1,\end{aligned}$$

where κ , τ and σ are called the first, the second and the third curvatures of the curve γ , respectively, [6].

$$\langle \mathbf{T}, \mathbf{T} \rangle = \langle \mathbf{N}, \mathbf{N} \rangle = \langle \mathbf{B}_1, \mathbf{B}_1 \rangle = \langle \mathbf{B}_2, \mathbf{B}_2 \rangle = 1.$$

The Bishop frame or parallel transport frame is an alternative approach to defining a moving frame that is well defined even when the curve has vanishing second derivative. One can express parallel transport of an orthonormal frame along a curve simply by parallel transporting each component of the frame, [4]. The tangent vector and any convenient arbitrary basis for the remainder of the frame are used. The Bishop frame is expressed as

$$\begin{aligned}\mathbf{T}' &= k_1\mathbf{M}_1 + k_2\mathbf{M}_2 + k_3\mathbf{M}_3, \\ \mathbf{M}'_1 &= -k_1\mathbf{T}, \\ \mathbf{M}'_2 &= -k_2\mathbf{T}, \\ \mathbf{M}'_3 &= -k_3\mathbf{T},\end{aligned}\tag{2.1}$$

where k_1, k_2, k_3 are principal curvature functions according to parallel transport frame of the curve γ and their expression as follows:

$$\begin{aligned}k_1 &= \kappa \cos \theta \cos \psi, \\ k_2 &= \kappa (-\cos \phi \sin \psi + \sin \phi \sin \theta \cos \psi), \\ k_3 &= \kappa (\sin \phi \sin \psi + \cos \phi \sin \theta \cos \psi),\end{aligned}$$

and

$$\begin{aligned}\kappa &= \sqrt{k_1^2 + k_2^2 + k_3^2}, \\ \tau &= -\psi' + \phi' \sin \theta, \\ \sigma &= \frac{\theta'}{\sin \psi}, \\ \phi' \cos \theta + \theta' \cot \psi &= 0,\end{aligned}$$

where

$$\theta' = \frac{\sigma}{\sqrt{\kappa^2 + \tau^2}}, \psi' = -\left(\tau + \sigma \frac{\sqrt{\sigma^2 - (\theta')^2}}{\sqrt{\kappa^2 + \tau^2}}\right), \phi' = \frac{\sqrt{\sigma^2 - (\theta')^2}}{\cos \theta}$$

and κ, τ, σ are the principal curvature functions according to Frenet frame of the curve γ , [3].

We shall call the set $\{\mathbf{T}, \mathbf{M}_1, \mathbf{M}_2, \mathbf{M}_3\}$ as parallel transport frame of γ .

3. Biharmonic curves according to parallel transport frame in \mathbb{E}^4

Biharmonic equation for the curve γ reduces to

$$\nabla_{\mathbf{T}}^3 \mathbf{T} - R(\mathbf{T}, \nabla_{\mathbf{T}} \mathbf{T}) \mathbf{T} = 0, \quad (3.1)$$

that is, γ is called a biharmonic curve if it is a solution of the equation (3.1).

Theorem 3.1. $\gamma : I \rightarrow \mathbb{E}^4$ is a unit speed biharmonic curve if and only if

$$\begin{aligned} k_1^2 + k_2^2 + k_3^2 &= \mathfrak{A}, \\ k_1'' - k_1(k_1^2 + k_2^2 + k_3^2) &= 0, \\ k_2'' - k_2(k_1^2 + k_2^2 + k_3^2) &= 0, \\ k_3'' - k_3(k_1^2 + k_2^2 + k_3^2) &= 0, \end{aligned} \quad (3.2)$$

where \mathfrak{A} is non-zero constant of integration.

Proof: Using the Bishop equations (2.1) and biharmonic equation (3.1), we obtain

$$\begin{aligned} (-3k_1'k_1 - 3k_2'k_2 - 3k_3'k_3)\mathbf{T} + (k_1'' - k_1(k_1^2 + k_2^2 + k_3^2))\mathbf{M}_1 \\ + (k_2'' - k_2(k_1^2 + k_2^2 + k_3^2))\mathbf{M}_2 - R(\mathbf{T}, \nabla_{\mathbf{T}} \mathbf{T}) \mathbf{T} = 0. \end{aligned} \quad (3.3)$$

In \mathbb{E}^4 , the Riemannian curvature is zero, we have

$$\begin{aligned} (-3k_1'k_1 - 3k_2'k_2 - 3k_3'k_3)\mathbf{T} + (k_1'' - k_1(k_1^2 + k_2^2 + k_3^2))\mathbf{M}_1 \\ + (k_2'' - k_2(k_1^2 + k_2^2 + k_3^2))\mathbf{M}_2 = 0. \end{aligned} \quad (3.4)$$

By (3.4), we see that γ is a unit speed biharmonic curve if and only if

$$\begin{aligned} -3k_1'k_1 - 3k_2'k_2 - 3k_3'k_3 &= 0, \\ k_1'' - k_1(k_1^2 + k_2^2 + k_3^2) &= 0, \\ k_2'' - k_2(k_1^2 + k_2^2 + k_3^2) &= 0, \\ k_3'' - k_3(k_1^2 + k_2^2 + k_3^2) &= 0. \end{aligned} \quad (3.5)$$

These, together with (3.5), complete the proof of the theorem. \square

Corollary 3.2. $\gamma : I \rightarrow \mathbb{E}^4$ is a unit speed biharmonic curve if and only if

$$\begin{aligned} k_1^2 + k_2^2 + k_3^2 &= \mathfrak{A}, \\ k_1'' - k_1\mathfrak{A} &= 0, \\ k_2'' - k_2\mathfrak{A} &= 0, \\ k_3'' - k_3\mathfrak{A} &= 0, \end{aligned} \quad (3.6)$$

where \mathfrak{A} is constant of integration.

Theorem 3.3. *Let $\gamma : I \rightarrow \mathbb{E}^4$ is a unit speed biharmonic curve, then*

$$\begin{aligned} k_1^2(s) + k_2^2(s) + k_3^2(s) &= \mathfrak{A}, \\ k_1(s) &= c_1 e^{\sqrt{\mathfrak{A}}s} + c_2 e^{-\sqrt{\mathfrak{A}}s}, \\ k_2(s) &= c_3 e^{\sqrt{\mathfrak{A}}s} + c_4 e^{-\sqrt{\mathfrak{A}}s}, \\ k_3(s) &= c_5 e^{\sqrt{\mathfrak{A}}s} + c_6 e^{-\sqrt{\mathfrak{A}}s}, \end{aligned} \tag{3.7}$$

where \mathfrak{A} , c_1 , c_2 , c_3 , c_4 , c_5 , c_6 are constants of integration.

Proof: Using (3.6), we have (3.7). □

Corollary 3.4. *If $c_1 = c_3 = c_5$ and $c_2 = c_4 = c_6$, then*

$$k_1(s) = k_2(s) = k_3(s).$$

Corollary 3.5. *Let $\gamma : I \rightarrow \mathbb{E}^4$ is a unit speed biharmonic curve, then*

$$\begin{aligned} c_1 e^{\sqrt{\mathfrak{A}}s} + c_2 e^{-\sqrt{\mathfrak{A}}s} &= \sqrt{\mathfrak{A}} \sin \Lambda \cos \Pi, \\ c_3 e^{\sqrt{\mathfrak{A}}s} + c_4 e^{-\sqrt{\mathfrak{A}}s} &= \sqrt{\mathfrak{A}} \sin \Lambda \sin \Pi, \\ c_5 e^{\sqrt{\mathfrak{A}}s} + c_6 e^{-\sqrt{\mathfrak{A}}s} &= \sqrt{\mathfrak{A}} \cos \Lambda, \end{aligned}$$

where $\Lambda \in [0, \pi]$, $\Pi \in [0, 2\pi)$ and \mathfrak{A} , c_1 , c_2 , c_3 , c_4 , c_5 , c_6 are constants of integration.

References

1. L.R. Bishop: *There is more than one way to frame a curve*, Amer. Math. Monthly, Volume 82, Issue 3, (1975) 246-251.
2. B. Bükcü, M.K. Karacan: *Bishop motion and Bishop Darboux rotation axis of the timelike curve in Minkowski 3-space*, Kochi J. Math. 4 (2009) 109–117.
3. G.Y. Jiang: *2-harmonic isometric immersions between Riemannian manifolds*, Chinese Ann. Math. Ser. A 7 (1986), 130–144.
4. F. Gökçelik, Z. Bozkurt, I. Gök, F. N. Ekmekci, Y. Yaylı: *Parallel transport frame in 4-dimensional Euclidean space E^4* , (preprint)
5. J. Milnor: *Curvatures of Left-Invariant Metrics on Lie Groups*, Advances in Mathematics 21 (1976), 293-329.
6. B. O'Neill: *Semi-Riemannian Geometry*, Academic Press, New York (1983).
7. C. Oniciuc: *On the second variation formula for biharmonic maps to a sphere*, Publ. Math. Debrecen 61 (2002), 613–622.
8. E. Turhan: *Completeness of Lorentz Metric on 3-Dimensional Heisenberg Group*, International Mathematical Forum, 13 (3) (2008), 639 - 644.
9. E. Turhan, T. Körpınar: *Characterize on the Heisenberg Group with left invariant Lorentzian metric*, Demonstratio Mathematica 42 (2) (2009), 423-428.
10. E. Turhan, T. Körpınar: *On Characterization Of Timelike Horizontal Biharmonic Curves In The Lorentzian Heisenberg Group $Heis^3$* , Zeitschrift für Naturforschung A- A Journal of Physical Sciences 65a (2010), 641-648.

11. E. Turhan and T. Körpınar: *Parametric equations of general helices in the sol space $\mathbb{S}oI^3$* , Bol. Soc. Paran. Mat. 31 (1) (2013), 99–104.
12. M. Turgut and S. Yılmaz: *Smarandache Curves in Minkowski Space-time*, International Journal of Mathematical Combinatorics 3 (2008), 51-55.

Talat Körpınar
Fırat University,
Department of Mathematics,
23119 Elazığ, Turkey
E-mail address: talatkorpınar@gmail.com

and

Essin Turhan
Fırat University,
Department of Mathematics,
23119 Elazığ, Turkey
E-mail address: essin.turhan@gmail.com