



On Fully-Convex Harmonic Functions and their Extension

Shahpour Nosrati and Ahmad Zireh

ABSTRACT: Uniformly convex univalent functions that introduced by Goodman, maps every circular arc contained in the open unit disk with center in it into a convex curve. On the other hand, a fully-convex harmonic function, maps each subdisk $|z| = r < 1$ onto a convex curve. Here we synthesis these two ideas and introduce a family of univalent harmonic functions which are fully-convex and uniformly convex also. In the following we will mention some examples of this subclass and obtain a necessary and sufficient conditions and finally a coefficient condition is given as an application of some convolution results.

Key Words: Uniformly convex function, Fully-Convex function, Harmonic function, Convolution.

Contents

1 Introduction and Preliminaries	51
2 Definition and Examples	54
3 Convolution and a sufficient condition	57

1. Introduction and Preliminaries

Let $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ be the open unit disk in complex plane. Let \mathcal{A} be the familier class of all analytic functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1.1)$$

in the open unit disk \mathbb{D} . Let \mathcal{S} denotes the family of all functions $f(z)$ of the form (1.1) that are univalent in \mathbb{D} and normalized with $f(0) = 0$ and $f'(0) = 1$.

A conformal function $f(z)$ is said to be starlike if every point of its range can be connected to the origin by a radial line that lies entirely in that region. The class of all starlike functions in \mathcal{S} is shown by \mathcal{S}^* [9] and $f(z) \in \mathcal{S}^*$ if and only if $\operatorname{Re} \left\{ z \frac{f'(z)}{f(z)} \right\} > 0$. Starlikeness is a hereditary property for conformal mappings, so if $f(z) \in \mathcal{S}$, and if f maps \mathbb{D} onto a domain that is starlike with respect to the origin, then the image of every subdisk $|z| < r < 1$ is also starlike with respect to the origin.

2010 *Mathematics Subject Classification*: Primary 30C45; Secondary 31C05, 31A05.
 Submitted December 29, 2016. Published October 04, 2017

An analytic function $f(z)$ is said to be convex if its range $f(\mathbb{D})$ is a convex set. It has shown that every convex function f in \mathcal{S} satisfy following analytic property

$$\operatorname{Re} \left\{ 1 + z \frac{f''(z)}{f'(z)} \right\} > 0$$

The class of all convex functions in \mathcal{S} is denoted by \mathcal{K} [9].

The subclass of uniformly starlike functions, \mathcal{UST} introduced by Goodman [6] and studied in analytic and geometric view.

Definition 1.1. [6] *A function $f(z) \in \mathcal{S}^*$ is said to be uniformly starlike in \mathbb{D} if it has the property that for every circular arc γ contained in \mathbb{D} , with center $\zeta \in \mathbb{D}$, the arc $f(\gamma)$ be starlike with respect to $f(\zeta)$. We denote the family of all uniformly starlike functions by \mathcal{UST} and we have,*

$$\mathcal{UST} = \left\{ f(z) \in \mathcal{S} : \operatorname{Re} \frac{(z - \zeta)f'(z)}{f(z) - f(\zeta)} > 0, (z, \zeta) \in \mathbb{D}^2 \right\} \quad (1.2)$$

It's clear that $\mathcal{UST} \subset \mathcal{S}^*$ and every function in \mathcal{UST} maps each subdisk $\{|z - \zeta| < \rho\} \subset \mathbb{D}$ onto a domain starlike with respect to $f(\zeta)$. Goodman [5] also defined the subclass of convex functions with this property that map each disk $\{|z - \zeta| < \rho\} \subset \mathbb{D}$ onto a convex domain and called it uniformly convex function and denoted the set of all these functions by \mathcal{UCV} :

Definition 1.2. [5] *A function $f(z) \in \mathcal{K}$ is said to be uniformly convex in \mathbb{D} if it has the property that for every circular arc γ contained in \mathbb{D} , with center $\zeta \in \mathbb{D}$, the arc $f(\gamma)$ be a convex arc. We have,*

$$\mathcal{UCV} = \left\{ f(z) \in \mathcal{S} : \operatorname{Re} \left(1 + (z - \zeta) \frac{f''(z)}{f'(z)} \right) \geq 0, (z, \zeta) \in \mathbb{D}^2 \right\} \quad (1.3)$$

A summary of early works on uniformly starlike and uniformly convex functions can be found in [10].

The complex-valued function $f(x, y) = u(x, y) + iv(x, y)$ is complex-valued harmonic function in \mathbb{D} if f is continuous and u and v are real harmonic in \mathbb{D} . We denote H the family of continuous complex-valued functions which are harmonic in the open unit disk \mathbb{D} . In simply-connected domain \mathbb{D} , $f \in H$ has a canonical representation $f = h + \bar{g}$, where h and g are analytic in \mathbb{D} [3,4]. Then, g and h have expansions in Taylor series as $h(z) = \sum_{n=0}^{\infty} a_n z^n$ and $g(z) = \sum_{n=0}^{\infty} b_n z^n$, so we may represent f by a power series of the form

$$f(z) = h(z) + \overline{g(z)} = \sum_{n=0}^{\infty} a_n z^n + \overline{\sum_{n=0}^{\infty} b_n z^n} \quad (1.4)$$

The Jacobian of a function $f = u + iv$ is $J_f(z) = \begin{vmatrix} u_x & u_y \\ v_x & v_y \end{vmatrix} = |h'(z)|^2 - |g'(z)|^2$, and $f(z) = h(z) + \overline{g(z)}$ is sense-preserving if $J_f(z) > 0$. In 1984, Clunie and Sheil-Small

[3] investigated the class S_H , consisting of sense-preserving univalent harmonic functions $f(z) = h(z) + \overline{g(z)}$ in simply-connected domain \mathbb{D} which normalized by $f(0) = 0$ and $f_z(0) = 1$ with the form,

$$f(z) = h(z) + \overline{g(z)} = z + \sum_{n=2}^{\infty} a_n z^n + \overline{\sum_{n=1}^{\infty} b_n z^n} \tag{1.5}$$

The subclass S_H^0 of S_H includes all functions $f \in S_H$ with $f_{\overline{z}}(0) = 0$, so $S \subset S_H^0 \subset S_H$. Clunie and Sheil-Small also considered convex functions in S_H , denoted by \mathcal{K}_H . The hereditary property of convexity for conformal maps does not generalize to univalent harmonic mappings. If f is a univalent harmonic map of \mathbb{D} onto a convex domain, then the image of the disk $|z| < r$ is convex for each radius $r \leq \sqrt{2} - 1$, but not necessarily for any radius in the interval $\sqrt{2} - 1 < r < 1$. In fact, the function

$$\begin{aligned} f(z) &= \mathbf{Re} \frac{z}{1-z} + i \mathbf{Im} \frac{z}{(1-z)^2} \\ &= \frac{z - \frac{1}{2}z^2}{(1-z)^2} + \frac{-\frac{1}{2}\overline{z}^2}{(1-\overline{z})^2} \in \mathcal{K}_H \end{aligned} \tag{1.6}$$

is a harmonic mapping of the disk \mathbb{D} onto the half-plane $\mathbf{Re} w > -\frac{1}{2}$, but the image of the disk $|z| \leq r$ fails to be convex for every r in the interval $\sqrt{2} - 1 < r < 1$ [4]. Thus we need a property to explain convexity of a map in a hereditary form in whole disk. We have following definition.

Definition 1.3. [2] *A harmonic mapping f with $f(0) = 0$ of the unit disk is said to be fully-convex if it maps every circle $|z| = r < 1$ in a one-to-one manner onto a convex curve.*

For $f \in S_H$, the family of fully-convex harmonic functions denotes by \mathcal{FK}_H . In 1980 Mocanu gave a relation between fully-starlikeness and a differential operator of a non-analytic function [7]. Let

$$Df = z f_z - \overline{z} f_{\overline{z}} \tag{1.7}$$

be the differential operator and

$$D^2 f = D(Df) = z z f_{zz} + \overline{z} \overline{z} f_{\overline{z}\overline{z}} + z f_z + \overline{z} f_{\overline{z}} \tag{1.8}$$

Lemma 1.4. [7] *Let $f \in C^1(\mathbb{D})$ is a complex-valued function such that $f(0) = 0$, $f(z) \neq 0$ for all $z \in \mathbb{D} - \{0\}$, and $J_f(z) > 0$ in \mathbb{D} and $\mathbf{Re} \frac{Df(z)}{f(z)} > 0$ then f is univalent and fully-starlike in \mathbb{D} .*

Lemma 1.5. *Let $f \in C^2(\mathbb{D})$ is a complex-valued function such that $f(0) = 0$, $f(z) \neq 0$ for all $z \in \mathbb{D} - \{0\}$, and $J_f(z) > 0$ in \mathbb{D} and $\mathbf{Re} \frac{D^2 f(z)}{Df(z)} > 0$ then f is univalent and fully-convex in \mathbb{D} .*

Since for a sense-preserving complex-valued function $f(z)$, $Df \neq 0$, If $f(z) \in \mathcal{S}_H$ and satisfies condition such as $\operatorname{Re} \frac{Df(z)}{f(z)} > 0$ or $\operatorname{Re} \frac{D^2f(z)}{Df(z)} > 0$ for all $z \in \mathbb{D} - \{0\}$, then f maps every circle $0 < |z| = r < 1$ onto a simple closed curve [7]. However, a fully-starlike mapping need not be univalent [2], we restrict our discussion to \mathcal{S}_H .

2. Definition and Examples

For a harmonic function $f(z) = h(z) + \overline{g(z)} \in \mathcal{S}_H$, and $\zeta \in \mathbb{D}$ we define the operator

$$\begin{aligned} \mathbf{D}f(z, \zeta) &= (z - \zeta)f_z(z) - \overline{(z - \zeta)}f_{\bar{z}}(z) \\ &= (z - \zeta)h'(z) - \overline{(z - \zeta)}g'(z) \end{aligned} \quad (2.1)$$

is harmonic also. For $\zeta = 0$ the operator $\mathbf{D}f(z, 0) = zf_z - \bar{z}f_{\bar{z}} = zh' - \bar{z}g' = Df(z)$ is previous operator (1.7). Differentiating of the operator $\mathbf{D}f(z, \zeta)$ gives us

$$\begin{aligned} \mathbf{D}^2f(z, \zeta) &= \mathbf{D}(\mathbf{D}f(z, \zeta)) \\ &= \mathbf{D}((z - \zeta)h'(z) - \overline{(z - \zeta)}g'(z)) \\ &= (z - \zeta)^2h''(z) + \overline{(z - \zeta)^2}g''(z) + (z - \zeta)h'(z) \\ &\quad + \overline{(z - \zeta)}g'(z) \end{aligned} \quad (2.2)$$

For $\zeta = 0$ the operator $\mathbf{D}^2f(z, 0) = z^2h''(z) + \bar{z}^2g''(z) + zh'(z) + \bar{z}g'(z) = D^2f(z)$ has described by Al-Amiri and Mocanu [1]. Similar to definition (1.1) we say that for an arbitrary function:

Definition 2.1. A function $f \in \mathcal{S}_H$ is said to be uniformly fully-convex harmonic function in \mathbb{D} if it has the property that for every circular arc γ contained in \mathbb{D} , with center $\zeta \in \mathbb{D}$, the arc $f(\gamma)$ is convex in $f(\mathbb{D})$.

We denote the set of all uniformly fully-convex harmonic functions in \mathbb{D} by \mathcal{UFK}_H . The following theorem gives analytic equivalency for above definition:

Theorem 2.2. Let $f \in \mathcal{S}_H$. $f \in \mathcal{UFK}_H$ if and only if

$$\operatorname{Re} \frac{D^2f(z, \zeta)}{Df(z, \zeta)} > 0, \quad (z, \zeta) \in \mathbb{D}^2 \quad (2.3)$$

Proof: Let $\gamma : \zeta + re^{i\theta}$ with $\theta_1 \leq \theta \leq \theta_2$ be a circular arc centered at ζ and contained in \mathbb{D} , then the image of γ under f is convex if the argument of the tangent to the image be a non-decreasing function of θ , that is,

$$\frac{\partial}{\partial \theta} \left(\arg \frac{\partial}{\partial \theta} \{f(z) - f(\zeta)\} \right) \geq 0$$

Hence

$$\operatorname{Im} \frac{\partial}{\partial \theta} \left(\log \frac{\partial}{\partial \theta} \{f(z) - f(\zeta)\} \right) \geq 0$$

But for a circular arc γ , set $z = \zeta + re^{i\theta}$, then $\frac{\partial}{\partial\theta}z = i(z - \zeta)$ and a brief computation will give us

$$\frac{\partial}{\partial\theta}\{f(z) - f(\zeta)\} = i\{(z - \zeta)f_z(z) - \overline{(z - \zeta)}f_{\bar{z}}(z)\} = i\mathbf{D}f(z, \zeta)$$

then

$$\begin{aligned} \frac{\partial}{\partial\theta} \log i\mathbf{D}f(z, \zeta) &= \frac{\partial}{\partial\theta} \log i\{(z - \zeta)h'(z) - \overline{(z - \zeta)}g'(z)\} \\ &= \frac{i[h'(z) + (z - \zeta)h''(z)]}{i\mathbf{D}f(z, \zeta)} i(z - \zeta) \\ &\quad - \frac{i[g'(z) + (z - \zeta)g''(z)]}{i\mathbf{D}f(z, \zeta)} \overline{i(z - \zeta)} \\ &= i \frac{\mathbf{D}^2 f(z, \zeta)}{\mathbf{D}f(z, \zeta)} \end{aligned}$$

Therefore, we must have

$$\mathbf{Im} \frac{\partial}{\partial\theta} \log i\mathbf{D}f(z, \zeta) = \mathbf{Re} \frac{\mathbf{D}^2 f(z, \zeta)}{\mathbf{D}f(z, \zeta)} \geq 0$$

as we want. \square

It should be noted that $\frac{\mathbf{D}^2 f(z, \zeta)}{\mathbf{D}f(z, \zeta)}(0, 0) = 1$, and

$$\mathcal{UFK}_H = \left\{ f(z) \in \mathcal{S}_H : \mathbf{Re} \frac{\mathbf{D}^2 f(z, \zeta)}{\mathbf{D}f(z, \zeta)} > 0, (z, \zeta) \in \mathbb{D}^2 \right\} \quad (2.4)$$

It's simple that one checks the rotations, $e^{-i\alpha}f(e^{i\alpha}z)$ for some real α , are preserve the class \mathcal{UFK}_H and the transformation $\frac{1}{t}f(tz)$ preserves this class also, where $0 < t \leq 1$. On the other hand, the class \mathcal{UFK}_H includes all fully-convex functions and uniformly convex functions. With $g = 0$ in (2.3), the analytic function $f(z) \in \mathcal{UFK}_H$ by (2.1) and (2.2) satisfies condition

$$\mathbf{Re} \frac{\mathbf{D}^2 f(z, \zeta)}{\mathbf{D}f(z, \zeta)} = \mathbf{Re} \frac{(z - \zeta)^2 h''(z) + (z - \zeta)h'(z)}{(z - \zeta)h'(z)} = \mathbf{Re} \left(1 + (z - \zeta) \frac{h''(z)}{h'(z)} \right) \geq 0$$

where $(z, \zeta) \in \mathbb{D}^2$. Then

Corollary 2.3. *If $f \in \mathcal{UCV}$ be an analytic function, then $f \in \mathcal{UFK}_H$. So, $\mathcal{UCV} \subset \mathcal{UFK}_H \subset \mathcal{K}_H$. Goodman [5] shows the analytic function $f(z) = \frac{z}{1 - Az} \in \mathcal{UCV}$ if and only if $|A| \leq \frac{1}{3}$, thus the convex function $f(z) = \frac{z}{1 - z} \notin \mathcal{UFK}_H$.*

Example 2.1. For $|\beta| < 1$ the affine mappings $f(z) = z + \overline{\beta z} \in \mathcal{UFK}_H$, since

$$\mathbf{Re} \frac{(z - \zeta) + \overline{(z - \zeta)\beta}}{(z - \zeta) - \overline{(z - \zeta)\beta}} \geq 0$$

is equivalent to

$$\mathbf{Re} \left((z - \zeta) + \overline{(z - \zeta)\beta} \right) \left(\overline{(z - \zeta)} - (z - \zeta)\beta \right) \geq 0$$

that is $(1 - |\beta|^2)|z - \zeta|^2 \geq 0$.

Corollary 2.4. For $\zeta = 0$ in (2.3), the harmonic function $f \in \mathcal{UFK}_H$ will be univalent and fully-convex in \mathbb{D} by Lemma 1.5. Thus it's clear any non fully-convex harmonic function is not in \mathcal{UFK}_H . The harmonic function $f(z) = \mathbf{Re} \frac{z}{1-z} + i\mathbf{Im} \frac{z}{(1-z)^2}$ isn't fully-convex ([4], p.46), then $f \notin \mathcal{UFK}_H$.

In the following we will give a necessary and sufficient condition for that $f \in \mathcal{UFK}_H$. This condition is a generalization form of a theorem about fully-convex functions mentioned by Chuaqui et al. in [2], p139.

Theorem 2.5. Let $f(z) \in \mathcal{S}_H$, $f \in \mathcal{UFK}_H$ if and only if

$$\begin{aligned} & |(z - \zeta)h'(z)|^2 \mathbf{Re} Q_h \geq \\ & |(z - \zeta)g'(z)|^2 \mathbf{Re} Q_g + \mathbf{Re} \left\{ (z - \zeta)^3 (h''(z)g'(z) - h'(z)g''(z)) \right\} \end{aligned} \quad (2.5)$$

where $Q_h = 1 + (z - \zeta) \frac{h''(z)}{h'(z)}$ and $Q_g = 1 + (z - \zeta) \frac{g''(z)}{g'(z)}$ for (z, ζ) in polydisk \mathbb{D}^2 .

Proof: According to the definition, $f \in \mathcal{UFK}_H$ if and only if $\mathbf{Re} \frac{\mathbf{D}^2 f(z, \zeta)}{\mathbf{D}f(z, \zeta)} > 0$ for $(z, \zeta) \in \mathbb{D}^2$, if and only if $\mathbf{Re} \left\{ \mathbf{D}^2 f(z, \zeta) \overline{\mathbf{D}f(z, \zeta)} \right\} > 0$ for $(z, \zeta) \in \mathbb{D}^2$, then a simple calculation gives us (2.5). \square

Lemma 2.6. $f = h + \overline{\beta h} \in \mathcal{UFK}_H$ if and only if $h \in \mathcal{UCV}$, where $|\beta| < 1$.

Proof: Let $f = h + \overline{g} \in \mathcal{S}_H$ and $g = \beta h$ with $|\beta| < 1$, then $f \in \mathcal{UFK}_H$ if and only if (2.5) holds. Since in this case, h and g satisfy equality $Q_h = Q_g$ so (2.5) holds if and only if $|(z - \zeta)h'(z)|^2 \mathbf{Re} Q_h (1 - |\beta|^2) \geq 0$, or $\mathbf{Re} Q_h \geq 0$ that shows $h \in \mathcal{UCV}$. \square

Example 2.2. The analytic function $h = z + Az^2$ is in \mathcal{UCV} if and only if $|A| \leq \frac{1}{6}$ [5]. By Lemma 2.6 we get $f(z) = z + Az^2 + \overline{\beta z + \beta Az^2} \in \mathcal{UFK}_H$ with $|\beta| < 1$ and $|A| \leq \frac{1}{6}$. For example, let $A = \frac{1}{6}$, $\beta = -\frac{i}{2}$ then $f = z + \frac{1}{6}z^2 - \frac{i}{2}z - \frac{i}{12}z^2 \in \mathcal{UFK}_H$. In Figure 1, the disk $|z - 0.7| < 0.3$ is mapped under this uniformly fully-convex harmonic function to a convex elliptical shape with center $f(\zeta) = (0.78, 0.39)$.




Figure 1: The image of $|z - 0.7| < 0.3$ under $f = z + \frac{1}{6}z^2 - \frac{i}{2}z - \frac{i}{12}z^2 \in \mathcal{UFK}_H$.

3. Convolution and a sufficient condition

The convolution or Hadamard product of two harmonic functions $f(z)$ and $F(z)$ with canonical representations

$$f(z) = h(z) + \overline{g(z)} = z + \sum_{n=2}^{\infty} a_n z^n + \sum_{n=1}^{\infty} \overline{b_n} \overline{z}^n \quad (3.1)$$

and

$$F(z) = H(z) + \overline{G(z)} = z + \sum_{n=2}^{\infty} A_n z^n + \sum_{n=1}^{\infty} \overline{B_n} \overline{z}^n \quad (3.2)$$

is defined as

$$(f * F)(z) = (h * H)(z) + \overline{g * G(z)} = z + \sum_{n=2}^{\infty} a_n A_n z^n + \sum_{n=1}^{\infty} \overline{b_n B_n} \overline{z}^n \quad (3.3)$$

The right half-plane mapping $\ell(z) = \frac{z}{1-z}$ acts as the convolution identity and the Koebe map $k(z) = \frac{z}{(1-z)^2}$ acts as derivative operation over functions convolution. We have some properties for convolution over analytic functions f and g :

$$\begin{aligned} f * g &= g * f & , & \quad \alpha(f * g) = \alpha f * g \\ f * \ell &= f & , & \quad z f'(z) = f * k(z) \end{aligned}$$

where $\alpha \in \mathbb{C}$. For a given subset $\mathcal{V} \subset \mathcal{A}$, its dual set \mathcal{V}^* is defined by

$$\mathcal{V}^* = \left\{ g \in \mathcal{A} : \frac{f * g(z)}{z} \neq 0, \forall f \in \mathcal{V}, \forall z \in \mathbb{D} \right\} \quad (3.4)$$

Nezhmetdinov (1997) proved that class \mathcal{UCV} is dual set for certain family of functions from \mathcal{A} . He proved ([8], Theorem 2, p.43) that the class \mathcal{UCV} is the dual set of a subset of \mathcal{A} consisting of functions $\varphi : \mathbb{D} \rightarrow \mathbb{C}$ given by

$$\varphi(z) = \frac{z}{(1-z)^3} \left[1 - z - \frac{4z}{(\alpha+i)^2} \right] \quad (3.5)$$

where $\alpha \in \mathbb{R}$. He determined the uniform estimate $|a_n(\varphi)| \leq n(2n-1)$ for the n -th Taylor coefficient of $\varphi(z)$:

Lemma 3.1. [8] *Let G is all function $\varphi \in \mathcal{A}$ of the form (3.5), then $\mathcal{UCV} = G^*$ and $|a_n(\varphi)| \leq n(2n-1)$ for all $n \geq 2$.*

For obtaining a sufficient condition in class \mathcal{UFK}_H , we define the dual set of a harmonic function. Let \mathcal{A}_H be the class of complex-valued harmonic functions $f(z) = h(z) + \overline{g(z)}$ in simply connected domain \mathbb{D} of the form (1.5) which are not necessarily sense-preserving univalent on \mathbb{D} . We define the dual set of a subset of \mathcal{A}_H :

Definition 3.2. *For a given subset $\mathcal{V}_H \subset \mathcal{A}_H$, the dual set \mathcal{V}_H^* is*

$$\mathcal{V}_H^* = \left\{ F = H + \overline{G} \in \mathcal{A}_H : \frac{h * H}{z} + \frac{\overline{g * G}}{\overline{z}} \neq 0, \forall f = h + \overline{g} \in \mathcal{V}_H, \forall z \in \mathbb{D} \right\} \quad (3.6)$$

Theorem 3.3. *Let $\alpha \in \mathbb{R}$, $|w| = 1$ and*

$$G_H = \left\{ \varphi - \sigma \overline{\varphi} : \varphi(z) = \frac{z}{(1-z)^3} \left(1 - \frac{w - i\alpha}{2 - w - i\alpha} z \right), \right. \\ \left. \sigma = \frac{\overline{(1-w)(2-w-i\alpha)}}{(1-w)(2-w-i\alpha)}, z \in \mathbb{D} \right\}$$

then $\mathcal{UFK}_H = G_H^*$. Furthermore If $\sum_{n=2}^{\infty} n(2n-1)|a_n| + n(2n-1)|b_n| < 1 - |b_1|$ then $f \in \mathcal{UFK}_H$.

It's clear that the analytic function φ is the same (3.5), but σ with $|\sigma| = 1$ isn't an arbitrary number and depend on both w and α in φ .

Proof: Let $f = h + \overline{g} \in \mathcal{UFK}_H$, that is

$$\operatorname{Re} \frac{(z-\zeta)^2 h''(z) + \overline{(z-\zeta)^2 g''(z)} + (z-\zeta)h'(z) + \overline{(z-\zeta)g'(z)}}{(z-\zeta)h'(z) - \overline{(z-\zeta)g'(z)}} > 0, \quad (3.7)$$

$(z, \zeta) \in \mathbb{D}^2$. For $\zeta = 0$ and then $z = 0$ we have $\frac{\mathbf{D}^2 f(z, \zeta)}{\mathbf{D}f(z, \zeta)} = 1$, hence the condition

(3.7) may be write as

$$i\alpha \left((z-\zeta)h'(z) - \overline{(z-\zeta)g'(z)} \right) \neq (z-\zeta)^2 h''(z) + \overline{(z-\zeta)^2 g''(z)} \\ + (z-\zeta)h'(z) + \overline{(z-\zeta)g'(z)}$$

where $\alpha \in \mathbb{R}$. By the minimum principle for harmonic functions, it is sufficient to verify this condition for $|z| = |\zeta|$ and so, we may assume that $\zeta = wz$ with $|w| = 1$, then from the definition of the dual set for harmonic functions (3.6), with straightforward calculation we conclude that $\frac{h * \varphi}{z} + \sigma \frac{\overline{g * \varphi}}{\bar{z}} \neq 0$, so the first assertion follows.

For obtaining coefficients condition, let $f(z) = h(z) + \overline{g(z)}$ is of the form (3.1), and $\varphi(z) = z + \sum_{n=2}^{\infty} \phi_n z^n$ be the series expansion of analytic function $\varphi(z)$, then $|\phi_n| \leq n(2n-1)$ for all $n \geq 2$, by Lemma 3.1. From previous part we see that

$$\begin{aligned} \left| \frac{h * \varphi}{z} + \sigma \frac{\overline{g * \varphi}}{\bar{z}} \right| &= \left| 1 + \sum_{n=2}^{\infty} a_n \phi_n z^{n-1} + \sigma \left(b_1 + \sum_{n=2}^{\infty} \overline{b_n \phi_n} \bar{z}^{n-1} \right) \right| \\ &\geq |1 + \sigma b_1| - \sum_{n=2}^{\infty} |a_n| |\phi_n| |z|^{n-1} - |\sigma| \sum_{n=2}^{\infty} |b_n| |\phi_n| |z|^{n-1} \\ &\geq |1 + \sigma b_1| - \sum_{n=2}^{\infty} n(2n-1) |a_n| - \sum_{n=2}^{\infty} n(2n-1) |b_n| \\ &> 0 \end{aligned}$$

when $\sum_{n=2}^{\infty} n(2n-1) |a_n| + n(2n-1) |b_n| < 1 - |b_1|$. □

References

1. Al-Amiri, H. and Mocanu, P. T., *Spirallike nonanalytic functions*, Proc. Amer. Math. Soc. **82** (1), 61-65, (1981).
2. Chuaqui, M., Duren, P. and Osgood, B., *Curvature properties of planar harmonic mappings*, Comput. Methods Funct. Theory **4** (1), 127-142, (2004).
3. Clunie, J. and Sheil-Small, T., *Harmonic Univalent Functions*, Ann. Acad. Sci. Fenn. Ser. A. I. Math. **9** (2), 3-25, (1984).
4. Duren, P. L., *Harmonic Mappings in the Plane*, Cambridge University Press, New York, (2004).
5. Goodman, A. W., *On Uniformly Convex Functions*, Ann. Polon. Math., **56**, 87-92, (1991).
6. Goodman, A. W., *On Uniformly Starlike Functions*, J. Math. Ana. & App. **155**, 364-370, (1991).
7. Mocanu, P. T., *Starlikeness and convexity for nonanalytic functions in the unit disc*, Mathematica (Cluj) **22** (45), 77-83, (1980).
8. Nezhmetdinov, I. R., *Classes of Uniformly Convex and Uniformly Starlike Functions as Dual Sets*, J. Math. Anal. Appl. **216**, 40-47, (1997).
9. Pommerenke, Ch., *Univalent Functions*, Vandenhoeck and Ruprecht, Göttingen, (1975).
10. Ronning, F., *A survey on uniformly convex and uniformly starlike functions*. Ann. Univ. Mariae Curie-Sklodowska Sect. A, **47** (13), 123-134, (1993).
11. Sheil-Small, T., *Constants for Planar Harmonic Mappings*, J. London Math. Soc. **2** (42), 237-248, (1990).

Shahpour Nosrati,
Faculty of Mathematical Sciences,
Shahrood University of Technology,
Iran.
E-mail address: shahpournosrati@yahoo.com

and

Ahmad Zireh,
Faculty of Mathematical Sciences,
Shahrood University of Technology,
Iran.
E-mail address: azireh@gmail.com