



Stochastic Stability and Impulsive Vaccination of Multicompartment Nonlinear Epidemic Model with Incidence Rate

Laid Chaharazed

ABSTRACT: In this work, we consider a multicompartment nonlinear epidemic model with temporary immunity and a saturated incidence rate. $N(t)$ at time t , this population is divide into seven sub-classes.

$N(t) = S(t) + E(t) + I(t) + I_1(t) + I_2(t) + I_3(t) + Q(t)$. where $S(t), E(t), I(t), I_1(t), I_2(t), I_3(t)$ and $Q(t)$ denote the sizes of the population susceptible to disease, exposed, infectious members and quarantine members with the possibility of infection through temporary immunity, respectively. The stability of a disease-free status equilibrium and the existence of endemic equilibrium determined by the ratio called the basic reproductive number. The multicompartment non linear epidemic model with saturated rate has been studied the stochastic stability of the free disease equilibrium under certain conditions, and obtain the conditions of global attractivity of the infection.

Key Words: Basic reproduction number, endemic equilibrium, epidemic, model stability, saturated incidence, stochastic stability, vaccination.

Contents

1 Introduction	1
2 Equilibrium points	2
2.1 Theorem 1.	3
2.2 Theorem 2.	3
2.3 Theorem 3.	3
3 Stochastic stability of the free disease equilibrium	4
3.1 Theorem 4.	4
4 Impulsive vaccination	5
4.1 Lemma 1.	6
4.2 Lemma 2.	6
4.3 Global attractivity	6
4.4 Theorem 5.	7
5 Conclusion	10

1. Introduction

This paper considers the following nonlinear epidemic model with temporary immunity:

$$\left\{ \begin{array}{l} \dot{S}(t) = \lambda + \nu - (\mu + d) S(t) - \beta \frac{S(t)I(t)}{1+aI^2(t)} + \gamma Q(t), \\ \dot{E}(t) = \beta \frac{S(t)I(t)}{1+aI^2(t)} - (\mu_1 + d) E(t) - \beta e^{-\mu_1 \tau} S(t - \tau) I(t - \tau), \\ \dot{I}(t) = \beta e^{-\mu_1 \tau} S(t - \tau) I(t - \tau) - (\mu_2 + d) I(t) - \sum_{i=1}^3 \alpha_i I(t), \\ \dot{I}_1(t) = \alpha_1 I(t) - \delta_1 I_1(t) - (\mu_3 + d) I_1(t), \\ \dot{I}_2(t) = \alpha_2 I(t) - \delta_2 I_2(t) - (\mu_4 + d) I_2(t), \\ \dot{I}_3(t) = \alpha_3 I(t) - \delta_3 I_3(t) - (\mu_5 + d) I_3(t), \\ \dot{Q}(t) = \sum_{i=1}^3 \delta_i I_i(t) - (\mu_6 + d) Q(t) - \gamma Q(t). \end{array} \right. \quad (1.1)$$

Consider a population of size $N(t)$ at time t , this population is divide into seven sub-classes, with $N(t) = S(t) + E(t) + I(t) + I_1(t) + I_2(t) + I_3(t) + Q(t)$.

Where $S(t), E(t), I(t), I_1(t), I_2(t), I_3(t)$ and $Q(t)$ denote the sizes of the population susceptible to disease, exposed, infectious members and quarantine members with the possibility of infection through temporary immunity, respectively. The positive constants $\mu, \mu_1, \mu_2, \mu_3, \mu_4, \mu_5$, and μ_6 represent the death rates of susceptible, exposed, infectious and quarantine. Biologically, it is natural to assume that $\mu \leq \min \{\mu_1, \mu_2, \mu_3, \mu_4, \mu_5, \mu_6\}$. The positive constant d is natural mortality rate. The positive constants λ represent rate of incidence. The positive constant γ represent the recovery rate of infection. The positive constant β is the average numbers of contacts infective for S and I . ν the positive constant is the parameter of emigration. The positive constant $\alpha_1, \alpha_2, \alpha_3$ are the average numbers of contacts. The positive constants $\delta_1, \delta_2, \delta_3$, are the numbers of transfer to conversion of infected people to quarantine.

a is saturation constant and τ is the length of immunity period.

The form of incidence is $\beta \frac{S(t)I(t)}{1+aI^2(t)}$; which is saturated with the susceptible.

The initial condition of (1) given as:

$$\begin{cases} S(\eta) = \Phi_1(\eta), E(\eta) = \Phi_2(\eta), I(\eta) = \Phi_3(\eta), I_1(\eta) = \Phi_4(\eta), \\ I_2(\eta) = \Phi_5(\eta), I_3(\eta) = \Phi_6(\eta), Q(\eta) = \Phi_7(\eta), -\tau \leq \eta \leq 0. \end{cases} \quad (1.2)$$

Where $\Phi = (\Phi_1, \Phi_2, \Phi_3, \Phi_4, \Phi_5, \Phi_6, \Phi_7)^T \in \mathbb{C}$ such that

$$\begin{cases} S(\eta) = \Phi_1(0) \geq 0, E(\eta) = \Phi_2(0) \geq 0, I(\eta) = \Phi_3(0) \geq 0, I_1(\eta) = \Phi_4(0) \geq 0, \\ I_2(\eta) = \Phi_5(0) \geq 0, I_3(\eta) = \Phi_6(0) \geq 0, Q(\eta) = \Phi_7(0) \geq 0, -\tau \leq \eta \leq 0. \end{cases} \quad (1.3)$$

Let \mathbb{C} denote the Banach space $C([-\tau, 0], \mathbb{R}^7)$ of continuous functions mapping the interval $[-\tau, 0]$ into \mathbb{R}^7 . With a biological meaning, we further assume that $\Phi_i(\eta) = \Phi_i(0) \geq 0$, for $i = 1, 2, 3, 4, 5, 6, 7$.

We have,

$$\dot{N}(t) = \lambda + \nu - \mu S(t) - \mu_1 E(t) - \mu_2 I(t) - \mu_3 I_1(t) - \mu_4 I_2(t) - \mu_5 I_3(t) - \mu_6 Q(t) - dN(t) \quad (1.4)$$

The region Ω is positively invariant set of (1.1),

$$\left\{ \begin{array}{l} \Omega = (S(t), E(t), I(t), I_1(t), I_2(t), I_3(t), Q(t)) \in \mathbb{R}^7, \\ S(t) + E(t) + I(t) + I_1(t) + I_2(t) + I_3(t) + Q(t) \leq N < \frac{\lambda + \nu}{\mu + d} \end{array} \right\} \quad (1.5)$$

Hence, system (1.1), can be rewrite as:

$$\left\{ \begin{array}{l} \dot{S}(t) = \lambda + \nu - (\mu + d) S(t) - \beta \frac{S(t)I(t)}{1+aI^2(t)} + \gamma Q(t), \\ \dot{E}(t) = \beta \frac{S(t)I(t)}{1+aI^2(t)} - (\mu_1 + d) E(t) - \beta e^{-\mu_1 \tau} S(t - \tau) I(t - \tau), \\ \dot{I}(t) = \beta e^{-\mu_1 \tau} S(t - \tau) I(t - \tau) - \left(\mu_2 + d + \sum_{i=1}^3 \alpha_i \right) I(t), \\ \dot{I}_1(t) = \alpha_1 I(t) - (\mu_3 + d + \delta_1) I_1(t), \\ \dot{I}_2(t) = \alpha_2 I(t) - (\mu_4 + d + \delta_2) I_2(t), \\ \dot{I}_3(t) = \alpha_3 I(t) - (\mu_5 + d + \delta_3) I_3(t), \\ \dot{Q}(t) = \sum_{i=1}^3 \delta_i I_i(t) - (\mu_6 + d + \gamma) Q(t). \end{array} \right. \quad (1.6)$$

2. Equilibrium points

We calculate the points of equilibrium in the absence and presence of infection.

In the absence of infection, the system (1.6) has a disease-free equilibrium E_0 .

$$E_0 = \left(\hat{S}, \hat{E}, \hat{I}, \hat{I}_1, \hat{I}_2, \hat{I}_3, \hat{Q} \right)^T = \left(\frac{\lambda + \nu}{\mu + d}, 0, 0, 0, 0, 0, 0 \right)^T. \quad (2.1)$$

Define the quarantine reproduction number as:

$$R_0 = \frac{\beta e^{-\mu_1 \tau}}{\mu + d} \times \frac{(\lambda + \nu)}{\mu_2 + d + \sum_{i=1}^3 \alpha_i} \quad (2.2)$$

In the presence of infection, substituting in the system, Ω also contains a unique positive, endemic equilibrium

$$E_\tau^* = (S^*, E^*, I^*, I_1^*, I_2^*, I_3^*, Q^*)^T \quad (2.3)$$

$$\left\{ \begin{array}{l} S^* = \frac{\mu_2 + d + \sum_{i=1}^3 \alpha_i}{\beta e^{-\mu_1 \tau}}, \\ E^* = \frac{1}{\mu_1 + d} \left[\beta S^* I^* \left(\frac{1}{1 + a(I^*)^2} - e^{-\mu_1 \tau} \right) \right] \\ I^* = \frac{\beta S^* + \sqrt{\Delta}}{2m}, m = \lambda + \nu - (\mu + d) S^* + \gamma Q^*, \Delta = (\beta S^*)^2 + 4\alpha m \\ I_1^* = \frac{\alpha_1}{\mu_3 + d + \delta_1} I^*, \\ I_2^* = \frac{\alpha_2}{\mu_4 + d + \delta_2} I^*, \\ I_3^* = \frac{\alpha_3}{\mu_5 + d + \delta_3} I^*, \\ O^* = \frac{\sigma}{c_3} I^*, \\ Q^* = \frac{1}{\mu_6 + d + \gamma} \left[\frac{\alpha_1 \delta_1}{\mu_3 + d + \delta_1} + \frac{\alpha_2 \delta_2}{\mu_4 + d + \delta_2} + \frac{\alpha_3 \delta_3}{\mu_5 + d + \delta_3} \right] I^*, \end{array} \right.$$

2.1. Theorem 1.

Let $(S, E, I, I_1, I_2, I_3, Q)$ the solution of (1.6) wich is defined in $[0, \infty)$.

$$\limsup_{t \rightarrow \infty} N(t) \leq \frac{\lambda + \nu}{\mu + d}.$$

PROOF.

We have (1.4), then for

$$\dot{N} \leq \lambda + \nu - (\mu + d) N. \quad (2.4)$$

With integrating, we have for

$$t \in (0, t_1), N \leq \frac{\lambda + \nu}{\mu + d} - \left(1 - e^{-(\mu + d)t} \right), \quad (2.5)$$

Then $N \leq 2 \frac{\lambda + \nu - \nu}{\mu + d}$. The solution $(S, E, I, I_1, I_2, I_3, Q)$ bounded in $[0, T)$. For

$$t \in [0, \infty), N \leq \frac{\lambda + \nu}{\mu + d} - \left(1 - e^{-(\mu + d)t} \right). \quad (2.6)$$

Finally

$$\limsup_{t \rightarrow \infty} N(t) \leq \frac{\lambda + \nu}{\mu + d}.$$

2.2. Theorem 2.

The disease-free equilibrium E_0 of the system (1.6) is locally asymptotically stable if $R_0 < 1$.

2.3. Theorem 3.

If $R_0 > 1$, the system (1.6) has a unique, non-trivial equilibrium E_τ^* which is locally asymptotically stable.

3. Stochastic stability of the free disease equilibrium

We limit ourselves here to perturbing only the contact rate so we replace β by $\beta + ab(t)$, where $b(t)$ is white noise (Brownian motion). The system (1.6) transformed to the following Itô stochastic differential equations:

$$\left\{ \begin{array}{l} dS = \left[\lambda + \nu - (\mu + d) S(t) - \beta \frac{SI}{1+aI^2} + \gamma Q(t) \right] dt - a \frac{SI}{1+aI^2} db, \\ dE = \left[\beta \frac{SI}{1+aI^2} - (\mu_1 + d) E(t) - \beta e^{-\mu_1 \tau} S(t - \tau) I(t - \tau) \right] dt + a \frac{SI}{1+aI^2} db, \\ dI = \left[\beta e^{-\mu_1 \tau} S(t - \tau) I(t - \tau) - \left(\mu_2 + d + \sum_{i=1}^3 \alpha_i \right) I(t) \right] dt, \\ dI_1 = [\alpha_1 I - (\mu_3 + d + \delta_1) I_1] dt, \\ dI_2 = [\alpha_2 I - (\mu_4 + d + \delta_2) I_2] dt, \\ dI_3 = [\alpha_3 I - (\mu_5 + d + \delta_3) I_3] dt, \\ dQ = \left[\sum_{i=1}^3 \delta_i I_i - (\mu_6 + d + \gamma) Q \right] dt. \end{array} \right. \quad (3.1)$$

3.1. Theorem 4.

If $R_0 < 1$, $E(t)$ and $Q(t)$ are exponentially almost surely stable.

PROOF: Let w , with Itô's formula, we obtain

$$L = d \log(E + wQ) = \frac{\dot{E} + w\dot{Q}}{E + wQ} \quad (3.2)$$

$$L = \frac{1}{E + wQ} \left[+w \left[\sum_{i=1}^3 \delta_i I_i - (\mu_6 + d + \gamma) Q \right] - \frac{1}{2} \frac{a^2}{E + wQ} \left(\frac{SI}{1+aI^2} \right)^2 \right] dt + \frac{a}{E + wQ} \times \frac{SI}{1+aI^2} db, \quad (3.3)$$

We have

$$L \leq \frac{1}{E + wQ} \left[- \left[\beta \frac{\lambda + \nu}{\mu + d} \left(\frac{1}{1+aI^2} - e^{-\mu_1 \tau} \right) \right] I - [\mu_1 + d] E + (w\delta_1) I_1 + (w\delta_2) I_2 + (w\delta_3) I_3 - w(\mu_6 + d + \gamma) Q \right] dt + \frac{a}{E + wQ} \times \frac{SI}{1+aI^2} db, \quad (3.4)$$

Then

$$L \leq \frac{-1}{E + wQ} \left[[\mu_1 + d] E + w(\mu_6 + d + \gamma) Q \right] dt + \frac{a}{E + wQ} \times \frac{SI}{1+aI^2} db, \quad (3.5)$$

We suppose that

$$L_1 = \min \{ (\mu_1 + d), (\mu_6 + d + \gamma) \}. \quad (3.6)$$

Then

$$L \leq -L_1 dt + \frac{aSI}{E + wQ} db. \quad (3.7)$$

With integration, we obtain

$$\log(E + wQ) \leq -L_1 t + a \int_0^t \frac{S(v) I(v)}{E(v) + wQ(v)} db(v). \quad (3.8)$$

We have $\left(\frac{S(v) I(v)}{E(v) + wQ(v)} \right)^2$, is bounded.

So, $\lim_{t \rightarrow \infty} \frac{S(v) I(v)}{E(v) + wQ(v)} db(v) = 0$, almost surely.

The following form from Doob's martingale inequality combined with Itô isometry see [18].

$\limsup_{t \rightarrow \infty} \frac{1}{t} \log(E + wQ) \leq -L_1$, almost surely.

Finally

- $\limsup_{t \rightarrow \infty} \frac{1}{t} \log E \leq -L_1$, almost surely,
- $\limsup_{t \rightarrow \infty} \frac{1}{t} \log Q \leq -L_1$, almost surely.

4. Impulsive vaccination

The following mathematical model with saturation incidence:

$$\left\{ \begin{array}{l} \left\{ \begin{array}{l} \dot{S}(t) = \lambda + \nu - (\mu + d) S(t) - \beta \frac{S(t)I(t)}{1+aI^2(t)} + \gamma Q(t), \\ \dot{E}(t) = \beta \frac{S(t)I(t)}{1+aI^2(t)} - (\mu_1 + d) E(t) - \beta e^{-\mu_1 \tau} S(t-\tau) I(t-\tau), \\ \dot{I}(t) = \beta e^{-\mu_1 \tau} S(t-\tau) I(t-\tau) - \left(\mu_2 + d + \sum_{i=1}^3 \alpha_i \right) I(t), \\ \dot{I}_1(t) = \alpha_1 I(t) - (\mu_3 + d + \delta_1) I_1(t), \\ \dot{I}_2(t) = \alpha_2 I(t) - (\mu_4 + d + \delta_2) I_2(t), \\ \dot{I}_3(t) = \alpha_3 I(t) - (\mu_5 + d + \delta_3) I_3(t), \\ \dot{Q}(t) = \sum_{i=1}^3 \delta_i I_i(t) - (\mu_6 + d + \gamma) Q(t), \end{array} \right. \right\}, t \neq kT, k \in \mathbb{N}, \\ \left\{ \begin{array}{l} S(t^+) = (1 - \theta) S(t), \\ E(t^+) = E(t), \\ I(t^+) = I(t^+), \\ I_1(t^+) = I_1(t^+), \\ I_2(t^+) = I_2(t^+), \\ I_3(t^+) = I_3(t^+), \\ Q(t^+) = Q(t) + \theta S(t), \end{array} \right\}, t = kT, k \in \mathbb{N}, \end{array} \right. \quad (4.1)$$

With θ , is pulse vaccination rate, ($0 < \theta < 1$).

We simplify (4.1), and with (1.4) the following system is:

$$\left\{ \begin{array}{l} \left\{ \begin{array}{l} \dot{S}(t) = \lambda + \nu - (\mu + d) S(t) - \beta \frac{S(t)I(t)}{1+aI^2(t)} + \gamma Q(t), \\ \dot{I}(t) = \beta e^{-\mu_1 \tau} S(t-\tau) I(t-\tau) - \left(\mu_2 + d + \sum_{i=1}^3 \alpha_i \right) I(t), \\ \dot{I}_1(t) = \alpha_1 I(t) - (\mu_3 + d + \delta_1) I_1(t), \\ \dot{I}_2(t) = \alpha_2 I(t) - (\mu_4 + d + \delta_2) I_2(t), \\ \dot{I}_3(t) = \alpha_3 I(t) - (\mu_5 + d + \delta_3) I_3(t), \\ \dot{Q}(t) = \sum_{i=1}^3 \delta_i I_i(t) - (\mu_6 + d + \gamma) Q(t), \\ \dot{N} \approx \lambda + \nu - (\mu + d) N(t) \end{array} \right. \right\}, t \neq kT, k \in \mathbb{N}, \\ \left\{ \begin{array}{l} S(t^+) = (1 - \theta) S(t), \\ I(t^+) = I(t^+), \\ I_1(t^+) = I_1(t^+), \\ I_2(t^+) = I_2(t^+), \\ I_3(t^+) = I_3(t^+), \\ Q(t^+) = Q(t) + \theta S(t), \\ N(t^+) = N(t), \end{array} \right\}, t = kT, k \in \mathbb{N}, \end{array} \right. \quad (4.2)$$

The initial condition of (4.2) given as:

$$\left\{ \begin{array}{l} S(\eta) = \Phi_1(\eta), I(\eta) = \Phi_2(\eta), I_1(\eta) = \Phi_3(\eta), \\ I_2(\eta) = \Phi_4(\eta), I_3(\eta) = \Phi_5(\eta), Q(\eta) = \Phi_6(\eta), N(\eta) = \Phi_7(\eta), -\tau \leq \eta \leq 0. \end{array} \right. \quad (4.3)$$

With a biological meaning, we further assume that: $\Phi_i(\eta) = \Phi_7(0) \geq 0$, for $i = 1, 2, 3, 4, 5, 6, 7$.

The region Ω_1 is positively invariant set of (4.2),

$$\left\{ \begin{array}{l} \Omega_1 = (S(t), I(t), I_1(t), I_2(t), I_3(t), Q(t), N(t)) \in \mathbb{R}^7, \\ S(t) + I(t) + I_1(t) + I_2(t) + I_3(t) + Q(t) \leq N < \frac{\lambda + \nu}{\mu + d}, N(t) \leq \frac{\lambda + \nu}{\mu + d} \end{array} \right\} \quad (4.4)$$

4.1. Lemma 1.

Consider the following impulsive differential system

$$\left\{ \begin{array}{l} \dot{u}(t) = a - bu(t), t \neq kT \\ u(t^+) = (1 - \theta)u(t), t = kT. \end{array} \right. \quad (4.5)$$

Where $a > 0, b > 0$, and $(0 < \theta < 1)$. Then there exists a unique positive periodic solution of system (4.2).

$$\tilde{u}_e(t) = \frac{a}{b} + \left(u^* - \frac{a}{b}\right) e^{-b(t-kT)}, kT < t \leq (k+1)T,$$

Which is globally asymptotically stable, where

$$u^* = \frac{a(1-\theta)(1-e^{-bT})}{b(1-(1-\theta)e^{-bT})}.$$

4.2. Lemma 2.

Consider the following linear neutral delay equation:

$$\dot{x}(t) = a_1x(t-\tau) - a_2x(t) \quad (4.6)$$

Where a_1, a_2 , and τ are all positive constants.

For $-\tau \leq t \leq 0, x(t) > 0$. We have:

$$If a_1 < a_2, \lim_{t \rightarrow \infty} x(t) = 0 \quad (4.7)$$

4.3. Global attractivity

In this section, we discuss the global attractivity of infection free periodic solution, in which infectious individuals are entirely absent from the population permanently

$$Q(\eta) = \Phi_\tau(0) \geq 0, -\tau \leq \eta \leq 0.$$

For all $t \geq 0$,

$$I(t) = 0, I_i(t) = 0.$$

Under this condition, the growth of susceptible individuals, quarantine individuals and total population must satisfy the following impulsive system:

$$\left\{ \begin{array}{l} \left\{ \begin{array}{l} \dot{S}(t) = \lambda + \nu - (\mu + d)S(t) + \gamma \frac{\lambda + \nu}{\mu + d} \frac{(1 - e^{-(\mu + d + \gamma)T})}{1 - (1 - \theta)e^{-(\mu + d + \gamma)T}} + \xi Q(t), \\ \dot{Q}(t) = -(\mu_6 + d + \gamma)Q(t), \\ \dot{N}(t) \approx \lambda + \nu - (\mu + d)N(t), \end{array} \right. \right\}, t \neq kT, k \in \mathbb{N}, \\ \left\{ \begin{array}{l} S(t^+) = (1 - \theta)S(t), \\ Q(t^+) = Q(t) + \theta S(t), \\ N(t^+) = N(t), \end{array} \right\}, t = kT, k \in \mathbb{N}, \end{array} \right. \quad (4.8)$$

We have

$$\left\{ \begin{array}{l} \dot{N}(t) = \lambda + \nu - (\mu + d)N(t), \\ N(t^+) = N(t), \end{array} \right. \quad (4.9)$$

From (4.9),

$$\lim_{t \rightarrow \infty} N(t) = \frac{\lambda + \nu}{\mu + d} \quad (4.10)$$

We have

$$\begin{cases} \dot{E}(t) = \beta \frac{S(t)I(t)}{1+aI^2(t)} - (\mu_1 + d)E(t) - \beta e^{-\mu_1 \tau} S(t-\tau)I(t-\tau), \\ E(t^+) = E(t), \end{cases} \quad (4.11)$$

From(4.11) and if $I(t) = 0$,

$$\lim_{t \rightarrow \infty} E(t) = 0 \quad (4.12)$$

We have S, Q oscillate with T in synchronization with the periodic pulse vaccination.

From (4.10) and (4.11), we have

$$Q(t) = \frac{\lambda + \nu}{\mu + d} - S(t), \quad (4.13)$$

and

$$\begin{cases} \dot{S}(t) = (\mu + d + \gamma) \left(\frac{\lambda + \nu}{\mu + d} - S(t) \right), t \neq kT \\ S(t^+) = (1 - \theta) S(t), t = kT \end{cases} \quad (4.14)$$

According to lemma 1, the periodic solution of system (4.14) is:

$$\tilde{S}_e(t) = \frac{\lambda + \nu}{\mu + d} + \left(S^* - \frac{\lambda + \nu}{\mu + d} \right) e^{-(\mu + d + \gamma)(t - kT)}, kT < t \leq (k + 1)T \quad (4.15)$$

It is globally asymptotically stable, with

$$S^* = \frac{\lambda + \nu}{\mu + d} \frac{(1 - \theta)(1 - e^{-(\mu + d + \gamma)T})}{1 - (1 - \theta)e^{-(\mu + d + \gamma)T}}. \quad (4.16)$$

4.4. Theorem 5.

The infection-free periodic solution $\left(\tilde{S}_e(t), 0, 0, 0, 0, \frac{\lambda + \nu}{\mu + d} - \tilde{S}_e(t), \frac{\lambda + \nu}{\mu + d} \right)$ of (4.2) is globally attractive provided that $Q^* < 1$.

$$Q^* = \frac{(\lambda + \nu) \beta e^{-\mu_1 \tau} (1 - e^{-(\mu + d + \gamma)T})}{(\mu + d) 1 - (1 - \theta) e^{-(\mu + d + \gamma)T}}.$$

PROOF.

With $Q^* < 1$, we choose ξ sufficiently small,

$$\beta e^{-\mu_1 \tau} \left(\frac{\lambda + \nu}{\mu + d} \frac{(1 - e^{-(\mu + d + \gamma)T})}{1 - (1 - \theta) e^{-(\mu + d + \gamma)T}} + \xi \right) < \mu_2 + d + \sum_{i=1}^3 \alpha_i \quad (4.17)$$

We have

$$\dot{S}(t) \leq (\lambda + \nu + \gamma) \left(\frac{\lambda + \nu}{\mu + d} - S(t) \right)$$

Then the following comparison impulsive differential system is:

$$\begin{cases} \dot{x}(t) = (\mu + d + \gamma) \left(\frac{\lambda + \nu}{\mu + d} - x(t) \right), t \neq kT \\ x(t^+) = (1 - \theta) x(t), t = kT \end{cases} \quad (4.18)$$

From (4.14), we have the periodic solution of (4.18) is (4.14), with $\tilde{x}_e(t) = \tilde{S}_e(t)$ which is globally asymptotically stable, if we have (4.15).

$(S(t), I(t), I_1(t), I_2(t), I_3(t), Q(t), N(t))$ the solution of (4.2), with (4.3) and $S(0^+) = S_0 > 0$ and $x(t)$ slution of (4.18), with $x(0^+) = S_0 > 0$.

By comparison theorem for impulsive differential system [12].
There exists an integer $k_1 > 0$, such that

$$S(t) < x(t) < \tilde{x}_e(t) + \xi, kT < t \leq (k+1)T, k > k_1, \quad (4.19)$$

From (4.19)

$$S(t) < \tilde{S}_e(t) + \xi \leq \left(\frac{\lambda + \nu}{\mu + d} \frac{(1 - e^{-(\mu+d+\gamma)T})}{1 - (1-\theta)e^{-(\mu+d+\gamma)T}} + \xi \right), kT < t \leq (k+1)T, k > k_1, \quad (4.20)$$

Then we have

$$S(t) \leq \sigma \quad (4.21)$$

With

$$\sigma = \frac{\lambda + \nu}{\mu + d} \frac{(1 - e^{-(\mu+d+\gamma)T})}{1 - (1-\theta)e^{-(\mu+d+\gamma)T}} + \xi.$$

From (4.21) and the second equation of (4.2), we have

$$\dot{I}(t) \leq \beta e^{-\mu_1 \tau} \sigma I(t - \tau) - \left(\mu_2 + d + \sum_{i=1}^3 \alpha_i \right) I(t), t > kT + \tau, k > k_1. \quad (4.22)$$

Consider the following comparison

$$\dot{y}(t) = \beta e^{-\mu_1 \tau} \sigma I(t - \tau) - \left(\mu_2 + d + \sum_{i=1}^3 \alpha_i \right) y(t), t > kT + \tau, k > k_1. \quad (4.23)$$

From (4.17),

$$\beta e^{-\mu_1 \tau} \sigma < \mu_2 + d + \sum_{i=1}^3 \alpha_i.$$

According to lemma 2,

$$\lim_{t \rightarrow \infty} y(t) = 0.$$

$y(t)$ is the solution of (4.23), with initial condition $y(\eta) = \Phi(\eta) \geq 0, -\tau \leq \eta \leq 0$.

We have $\lim_{t \rightarrow \infty} I(t) \leq \lim_{t \rightarrow \infty} y(t) = 0$, then

$$\lim_{t \rightarrow \infty} I(t) = 0. \quad (4.24)$$

For $\xi_1 > 0$ sufficiently small, there exist an integer $k_2 > 0$, with $k_2 T > kT + \tau$, with $I(t) < \xi_1$, for all $t > k_2 T$.

We have

$$\begin{cases} \dot{I}_1(t) = \alpha_1 I(t) - (\mu_3 + d + \delta_1) I_1(t), \\ I(t) < \xi_1, \text{ for all } t > k_2 T. \end{cases} \quad (4.25)$$

Then for $t > k_3 T$, we have

$$\dot{I}_1(t) < \alpha_1 \xi_1 - (\mu_3 + d + \delta_1) I_1(t) \quad (4.26)$$

Consider comparison equation for $t > k_2 T$

$$\dot{L}_1(t) = \alpha_1 \xi_1 - (\mu_3 + d + \delta_1) L_1(t) \quad (4.27)$$

We see $\lim_{t \rightarrow \infty} L_1(t) = 0$, by comparison theorem [12] there exists an integer $k_4 > k_3$, such that for all $t > k_3 T$, and ξ_1 arbitrarily small, we have

$$\lim_{t \rightarrow \infty} I_1(t) = 0 \quad (4.28)$$

We have

$$\begin{cases} \dot{I}_2(t) = \alpha_2 I(t) - (\mu_4 + d + \delta_2) I_2(t), \\ I(t) < \xi_1, \text{ for all } t > k_2 T. \end{cases} \quad (4.29)$$

Then for $t > k_4 T$, we have

$$\dot{I}_2(t) < \alpha_2 \xi_1 - (\mu_4 + d + \delta_2) I_2(t) \quad (4.30)$$

Consider comparison equation for $t > k_2 T$

$$\dot{L}_2(t) = \alpha_2 \xi_1 - (\mu_4 + d + \delta_2) L_2(t) \quad (4.31)$$

We see $\lim_{t \rightarrow \infty} L_2(t) = 0$, by comparison theorem [12] there exists an integer $k_5 > k_4$, such that for all $t > k_4 T$, and ξ_1 arbitrarily small, we have

$$\lim_{t \rightarrow \infty} I_2(t) = 0 \quad (4.32)$$

We have

$$\begin{cases} \dot{I}_3(t) = \alpha_3 I(t) - (\mu_5 + d + \delta_3) I_3(t), \\ I(t) < \xi_1, \text{ for all } t > k_2 T. \end{cases} \quad (4.33)$$

Then for $t > k_5 T$, we have

$$\dot{I}_3(t) < \alpha_3 \xi_1 - (\mu_5 + d + \delta_3) I_3(t) \quad (4.34)$$

Consider comparison equation for $t > k_2 T$

$$\dot{L}_3(t) = \alpha_3 \xi_1 - (\mu_5 + d + \delta_3) L_3(t) \quad (4.35)$$

We see $\lim_{t \rightarrow \infty} L_3(t) = 0$, by comparison theorem [12] there exists an integer $k_6 > k_5$, such that for all $t > k_5 T$, and ξ_1 arbitrarily small, we have

$$\lim_{t \rightarrow \infty} I_3(t) = 0 \quad (4.36)$$

From (4.10), and (4.24), there exists $k_7 > k_6$ such that

$$I(t) < \xi_1, \text{ and } N(t) > \frac{\lambda + \nu}{\mu + d} - \xi_1, \text{ for } t > k_7 T \quad (4.37)$$

From second equation of (4.1), Then

$$\dot{E}(t) \leq \frac{\lambda + \nu}{\mu + d} \frac{\beta \xi_1}{1 + a \xi_1^2} - (\mu_1 + d) E(t), \text{ for } t > k_7 T \quad (4.38)$$

There exists $k_8 > k_7$ such that

$$E(t) < \frac{\lambda + \nu}{(\mu + d)^2} \frac{\beta \xi_1}{1 + a \xi_1^2} + \xi_1, \text{ for } t > k_8 T \quad (4.39)$$

We have the first equation to (4.2), (4.37), and (4.39), we have

$$\begin{aligned} \dot{S}(t) &\geq \lambda + \nu - (\mu + d) S(t) - \beta \frac{S(t) \xi_1}{1 + a \xi_1^2(t)} + \gamma \left(\frac{\lambda + \nu}{\mu + d} - \frac{\beta \xi_1}{1 + a \xi_1^2} \frac{\lambda + \nu}{(\mu + d)^2} - S(t) - 3 \xi_1 \right) \\ \dot{S}(t) &\geq \lambda + \nu + \gamma \frac{\lambda + \nu}{\mu + d} - \frac{\gamma \beta \xi_1}{1 + a \xi_1^2} \frac{\lambda + \nu}{(\mu + d)^2} - 3 \gamma \xi_1 - \left(\mu + d + \frac{\beta \xi_1}{1 + a \xi_1^2(t)} + \gamma \right) S(t), \end{aligned} \quad (4.40)$$

Consider the comparison impulsive differential equation for $t > k_8 T$ and $k > k_8$,

$$\begin{cases} \dot{u}(t) = \left(\lambda + \nu + \gamma \frac{\lambda + \nu}{\mu + d} - \frac{\gamma \beta \xi_1}{1 + a \xi_1^2} \frac{\lambda + \nu}{(\mu + d)^2} - 3 \gamma \xi_1 \right) - \left(\mu + d + \frac{\beta \xi_1}{1 + a \xi_1^2(t)} + \gamma \right) u(t), t \neq kT \\ u(t^+) = (1 - \theta) u(t), t = kT \end{cases} \quad (4.41)$$

With lemma 1, we have the solution

$$\tilde{u}_e(t) = \Psi + (u^* - \Psi) e^{-\left(\mu+d+\frac{\beta\xi_1}{1+a\xi_1^2(t)}+\gamma\right)(t-kT)}, kT < t \leq (k+1)T,$$

Which is globally asymptotically stable, where

$$\Psi = \frac{\lambda + \nu + \gamma \frac{\lambda+\nu}{\mu+d} - \frac{\gamma\beta\xi_1}{1+a\xi_1^2} \frac{\lambda+\nu}{(\mu+d)^2} - 3\gamma\xi_1}{\mu + d + \frac{\beta\xi_1}{1+a\xi_1^2(t)} + \gamma}, u^* = \Psi \frac{(1-\theta) \left(1 - e^{-\left(\mu+d+\frac{\beta\xi_1}{1+a\xi_1^2(t)}+\gamma\right)T}\right)}{1 - (1-\theta) e^{-\left(\mu+d+\frac{\beta\xi_1}{1+a\xi_1^2(t)}+\gamma\right)T}}.$$

Consider the comparasion impulsive differential equation, there exists an integer $k_9 > k_8$,

$$S(t) > \tilde{u}_e(t) - \xi_1, kT < t \leq (k+1)T, k > k_9 \quad (4.42)$$

From (4.20), (4.42), we have

$$\tilde{S}_e(t) = \frac{\lambda + \nu}{\mu + d} \left(1 - \frac{\theta}{1 - (1-\theta) e^{-(\mu+d+\gamma)T}} e^{-(\mu+d+\gamma)(t-kT)}\right), kT < t \leq (k+1)T, \quad (4.43)$$

(4.43) is globally attractive, then

$$\lim_{t \rightarrow \infty} S(t) = \tilde{S}_e(t). \quad (4.44)$$

From (4.43), we have

$$\lim_{t \rightarrow \infty} E(t) = 0. \quad (4.45)$$

We have

$$\lim_{t \rightarrow \infty} Q(t) = \frac{\lambda + \nu}{\mu + d} - \tilde{S}_e(t), \quad (4.46)$$

From (4.10), (4.24), (4.28), (4.32), (4.36), (4.44), (4.45), and (4.46), the infection free periodic solution $\left(\tilde{S}_e(t), 0, 0, 0, 0, \frac{\lambda+\nu}{\mu+d} - \tilde{S}_e(t), \frac{\lambda+\nu}{\mu+d}\right)$ is globally attractive.

5. Conclusion

This paper addresses the equilibrium and stability of the epidemic model with temporary immunity and saturated incidence rate. Both trivial and endemic equilibrium are founds.

The disease-free equilibrium E_0 is globally asymptotically stable, if $R_0 < 1$, and the system has a unique non-trivial equilibrium E_T^* which is globally asymptotically stable if $R_0 > 1$. We study stochastic stability of the free disease equilibrium under some conditions, and finally we analyse impulsive vaccination. using theorem 5 imply that the disease dynamics of (22) is determined with $Q^* < 1$.

References

1. A. Lahrouz, L. Omari, D. Kiouach. Global analysis of a deterministic and stochastic nonlinear SIRS epidemic model. *Nonlinear Analysis: Modelling and Control*, Vol. 16, No. 1, pp 59–76, (2011).
2. Anderson R. M and Medley R M and Johnson. A. K. A Preliminary Study of the Transmission Dynamics of the Human Immunodeficiency Virus (HIV), the Causative Agent of AIDS. *IMA. J. Math. Appl. Med. Biol* 3, pp. 229-263, (1986).
3. Abta A and Kaddar A and Talibi H. A. Global Stability for Delay SIR and SEIR Epidemic Models with Saturated Incidence Rates. *Electronic Journal of Differential Equations*, 23, pp. 1-13, (2012).
4. Bailey. N.T.J-1. The Mathematical Theory of Infection Diseases and its Application. *Applied Statistics*, 26, N1, pp. 85-87, (1977).
5. Batiha, M. S. M. Noorani and I. Hashim. Numerical solutions of the nonlinear integro-differential equations, *Int. J. Open Probl. Compt. Math.* pp. 34-42, (2008).
6. Billard. L .A Stochastic General Epidemic in m Sub-Population. *J. Appl. Prob.* 13, pp. 567-572, (1976).
7. Dongmei Xio and Shigui Ruan. Global analysis of an epidemic model with non monotone incidence rate. *Mathematical Biosciences* 208, pp 419-429, (2007).

8. James M. Hyman, Jia Li. Epidemic Models with differential susceptibility and staged progression and their dynamics. *Mathematical Biosciences and engineering*. Volume 6, Number 2, April , pp. 321–332, (2009).
9. Hamid El Maroufy, A dil Lahrouz and PGL Leach. Qualitative Behaviour of a Model of an SIRS Epidemic. *Applied Mathematics & Information Sciences*. 5 (2) , pp 220-238, (2011).
10. Jinliang W, Xinxin Tian. Global Stability of a Delay Differential Equation of Hepatitis B Virus Infection with Immune Response. *Electronic Journal of Differential Equations*, 94, pp.1-11, (2013).
11. Jin. Z, Zhien. M and Maoan. H. Global stability of an SIRS epidemic model with delay. *Acta Matimatica Scientia*. 26. B, pp. 291-306, (2006).
12. Lakshmikantham, V., Bainov, D.D., Simeonov, P.S. *Theory of Impulsive Differential Equations*. World Scientific, Singapore.(1989).
13. Luo Q and Mao X. Stochastic population dynamics under regime switching. *J. Math. Anal. Appl.*334, pp. 69-84, (2007).
14. Naresh R and Omar S. An epidemic model for the transmission dynamics of HIV/AIDS and another infection. *International Journal of Mathematical Archive-1*(3, pp. 68-72), (2010).
15. P. LaSalle and S. Lefschetz. *Stability by Liapunov's direct method*. Academic Press. (1961).
16. Øksendal. B. *Stochastic Differential Equations an Introduction with Applications*, 5th ed. Springer, (2000).
17. Qin Zou, Shujing Gao, Qi Zhong. Pulse Vaccination Strategy in an Epidemic Model with Time Delays and Nonlinear Incidence. *Advanced Studies in Biology*, Vol. 1, no. 7, 307 - 321, (2009).
18. Ray Waston. On the Size Distribution for Some Epidemic Models. *J. Appl. Prob.*17, pp. 912-921, (1980).
19. Robert N and May. *Population Biology of infectious diseases I*. International centre of theoretical physics, pp. 1-9, (1982).
20. Ruoyan Sun. Global stability of the endemic equilibrium of multi group SIR models with nonlinear incidence. *Computers and Mathematics with Applications* 60, pp. 2286-2291, (2010).
21. Sarah A, Farida M, Muna A. Stability Analysis of an HIV/AIDS Epidemic. *International Mathematical Forum*. Vol 6, no 66.pp 3251-3273, (2011).
22. Shujing Gao,a, Lansun Chenc, Zhidong Tenga. Impulsive Vaccination of an SEIRS Model with Time Delay and Varying Total Population Size. *Bulletin of Mathematical Biology* 69: 731–745, (2007).
23. S. Pathak, A. Maiti, G.P. Samanta. Rich dynamics of an SIR epidemic model. *Nonlinear Analysis: Modelling and Control*, Vol. 15, No. 1, 71–81 2010.
24. S. Gao, L. Chen, J.J. Nieto, A. Torres, Analysis of a delayed epidemic model with pulse vaccination and saturation incidence, *Vaccine*, 24, 6037 - 6045, (2006).
25. Volodymyr Makarov, Denis Dragunov. A numeric-analytical method for solving the cauchy problem for ordinary differential equations. *Applied Mathematics and Computation*, pp. 1-26, (2010).
26. W. Ma, Y. Takeuchi, T. Hara and E. Beretta. Permanence of an SIR epidemic model with distributed time delays. *Tohoku Math. J.* 54, pp. 581-591, (2002).
27. W. Wang, Global. Behavior of an SEIR epidemic model with time delay. *Appl. Math. Letters*.15, pp. 423-428, (2002).
28. Wen L and Yang X. Global stability of a delayed SIRS model with temporary immunity. *Chaos, Solitons and Fractals* 38, pp. 221-226, (2008).
29. Xiao, L Chen and F. Ven den Bosch. Dynamical behavior for a stage-structured SIR infectious disease model. *Nonlinear Anal. Real World Appl* 3, pp.175-190, (2002).
30. Xia Wang, Youde Tao , Xinyu Song. Pulse vaccination on SEIR epidemic model with nonlinear incidence rate. *Applied Mathematics and Computation*. pp.398-404, (2009).
31. Y. Kuang. *Delay Differential Equation with Application in Population Dynamics*, Academic Press, New York, (1993).
32. Y. Xiao, L. Chen. Modelling and analysis of a predator-prey model with disease in the prey, *Math. Biosci*, 171, 59 - 82, (2001).
33. Z. Ma, J. Liu and J. Li. Stability analysis for differential infectivity epidemic models. *Nonlinear Anal. Real World Appl* 4, pp. 841-856, (2003).
34. Zhang F and Zhen Li and Zhang F. Global stability of an SIR epidemic model with constant infectious period. *Applied Mathematics and Computation* 199, pp.285-291, (2008).

Laid Chaharazed,
Department of Mathematics,
Faculty of Exact Sciences,
University Freres Mentouri,
Constantine 1, Algeria.
E-mail address: chahrazed2009@yahoo.fr; laid-chahrazed@umc.edu.dz